

Rabbit anti-Human ATG4C Polyclonal Antibody | anti-ATG4C antibody

Rabbit anti-ATG4C Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220447	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ATG4C; Polyclonal Antibody; Rabbit anti-ATG4C Antibody; APG4C; AUTL1; AUTL3; Cysteine protease ATG4C; AUT-like 3 cysteine endopeptidase; Autophagin-3; Autophagy-related cysteine endopeptidase 3; Autophagy-related protein 4 homolog C; anti-ATG4C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-ATG4C antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human ATG4C. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ATG4C staining in K562 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of ATG4C staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ATG4C expression in K562 (A) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-ATG4C antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	30410844
NCBI GeneID	84938
NCBI Accession #	NP_116241.2
NCBI GenBank Nucleotide #	NM_032852.3
UniProt Accession #	Q96DT6; Q96K04; A6NLR8; D3DQ58
Molecular Weight	52,497 Da
NCBI Official Full Name	cysteine protease ATG4C
NCBI Official Synonym Full Names	autophagy related 4C, cysteine peptidase
NCBI Official Symbol	ATG4C
NCBI Official Synonym Symbols	APG4C; AUTL1; AUTL3; APG4-C
NCBI Protein Information	cysteine protease ATG4C; autophagin-3; APG4 autophagy 4 homolog C; AUT-like 3 cysteine endopeptidase; ATG4 autophagy related 4 homolog C; AUT-like 1, cysteine endopeptidase; autophagy-related protein 4 homolog C; autophagy-related cysteine endopeptidase 3
UniProt Protein Name	Cysteine protease ATG4C
UniProt Gene Name	ATG4C
UniProt Synonym Gene Names	APG4C; AUTL1; AUTL3
UniProt Entry Name	ATG4C_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/atg4c-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220447>