

Rabbit anti-Human Ki67 Polyclonal Antibody | anti-Ki67 antibody

Rabbit anti-Ki67 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220612	Rabbit	Human	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

Ki67; Polyclonal Antibody; Rabbit anti-Ki67 Antibody; Antigen KI-67; anti-Ki67 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-Ki67 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Ki67. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of Ki67 staining in COS7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of Ki67 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of Ki67 expression in HCT116 (A), U87MG (B), HepG2 (C), LO2 (D), SKOVCA3 (E) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-Ki67 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	225543215
NCBI GeneID	4288
NCBI Accession #	NP_001139438.1
NCBI GenBank Nucleotide #	NM_001145966.1
UniProt Accession #	P46013; Q5VWH2
Molecular Weight	358,694 Da
NCBI Official Full Name	antigen KI-67 isoform 2
NCBI Official Synonym Full Names	antigen identified by monoclonal antibody Ki-67
NCBI Official Symbol	MKI67
NCBI Official Synonym Symbols	KIA
NCBI Protein Information	antigen KI-67; proliferation-related Ki-67 antigen
UniProt Protein Name	Antigen KI-67
UniProt Gene Name	MKI67
UniProt Entry Name	KI67_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ki67-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220612>