

Rabbit DGK alpha Polyclonal Antibody | anti-DGKalpha antibody

Rabbit anti-DGK alpha Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA220875	Rabbit	Human, Monkey	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

DGK alpha; Polyclonal Antibody; Rabbit anti-DGK alpha Antibody; DAGK; DAGK1; Diacylglycerol kinase alpha; DAG kinase alpha; 80 kDa diacylglycerol kinase; Diglyceride kinase alpha; DGK-alpha; anti-DGKalpha antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-DGKalpha antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human DGK alpha. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of DGK alpha staining in MDCK cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of DGK alpha staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of DGK alpha expression in MDCK (A), COS7 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-DGKalpha antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	31542504
NCBI GeneID	1606
NCBI Accession #	NP_001336.2
NCBI GenBank Nucleotide #	NM_001345.4
UniProt Accession #	P23743; O75481; O75482; O75483; O95217; Q8IZ56; Q8N5Q2
Molecular Weight	82,630 Da
NCBI Official Full Name	diacylglycerol kinase alpha
NCBI Official Synonym Full Names	diacylglycerol kinase, alpha 80kDa
NCBI Official Symbol	DGKA
NCBI Official Synonym Symbols	DAGK; DAGK1; DGK-alpha
NCBI Protein Information	diacylglycerol kinase alpha; DAG kinase alpha; diglyceride kinase alpha; 80 kDa diacylglycerol kinase
UniProt Protein Name	Diacylglycerol kinase alpha
UniProt Gene Name	DGKA
UniProt Synonym Gene Names	DAGK; DAGK1; DAG kinase alpha; DGK-alpha
UniProt Entry Name	DGKA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/dgk-alpha-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/220875>