

Rabbit CD137 Polyclonal Antibody | anti-CD137 antibody

Rabbit anti-CD137 Antibody

CATALOG # AAA221567	HOST Rabbit	REACTIVITY Human, Mouse, Monkey	APPLICATIONS Immunofluorescence, Western Blot
-------------------------------	-----------------------	---	---


PRODUCT SYNONYMS

CD137; Polyclonal Antibody; Rabbit anti-CD137 Antibody; ILA; Tumor necrosis factor receptor superfamily member 9; 4-1BB ligand receptor; CDw137; T-cell antigen 4-1BB homolog; T-cell antigen ILA; anti-CD137 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CD137 antibody	IC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD137. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of CD137 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



WB (Western Blot)

Western blot analysis of CD137 expression in HepG2 (A), NIH3T3 (B), mouse brain (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CD137 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	5730095
NCBI GeneID	3604
NCBI Accession #	NP_001552.2
NCBI GenBank Nucleotide #	NM_001561.5
UniProt Accession #	Q07011; P20334; AAB34924
Molecular Weight	27,899 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 9
NCBI Official Synonym Full Names	tumor necrosis factor receptor superfamily, member 9
NCBI Official Symbol	TNFRSF9
NCBI Official Synonym Symbols	ILA; 4-1BB; CD137; CDw137
NCBI Protein Information	tumor necrosis factor receptor superfamily member 9; CD137 antigen; T cell antigen ILA; T-cell antigen ILA; 4-1BB ligand receptor; homolog of mouse 4-1BB; receptor protein 4-1BB; T-cell antigen 4-1BB homolog; induced by lymphocyte activation (ILA); interl
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 9
UniProt Gene Name	TNFRSF9
UniProt Synonym Gene Names	CD137; ILA
UniProt Entry Name	TNR9_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/cd137-antibody-immunochemistry-immunofluorescence-western-blot-elisa/221567>