

Rabbit GRP75 Polyclonal Antibody | anti-GRP75 antibody

Rabbit anti-GRP75 Antibody

CATALOG # AAA222487	HOST Rabbit	REACTIVITY Human, Mouse, Rat	APPLICATIONS Immunofluorescence, Immunohistochemistry, Western Blot
-------------------------------	-----------------------	--	---


PRODUCT SYNONYMS

GRP75; Polyclonal Antibody; Rabbit anti-GRP75 Antibody; HSPA9; HSPA9B; Stress-70 protein; mitochondrial; 75 kDa glucose-regulated protein; GRP-75; Heat shock 70 kDa protein 9; Mortalin; MOT; Peptide-binding protein 74; PBP74; anti-GRP75 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-GRP75 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human GRP75. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of GRP75 staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of GRP75 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of GRP75 expression in HeLa (A), NIH3T3 (B), SP2/0 (C), mouse brain (D), rat brain (E) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-GRP75 antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	21264428
NCBI GeneID	3313
UniProt Accession #	P38646 ; Q8N1C8 ; P38647 ; P48721
Molecular Weight	679
NCBI Official Full Name	Stress-70 protein, mitochondrial
NCBI Official Synonym Full Names	heat shock 70kDa protein 9 (mortalin)
NCBI Official Symbol	HSPA9
NCBI Official Synonym Symbols	CSA; MOT; MOT2; CRP40; GRP75; PBP74; GRP-75; HSPA9B; MTHSP75; HEL-S-124m
NCBI Protein Information	stress-70 protein, mitochondrial; mortalin-2; p66-mortalin; mortalin, perinuclear; heat shock 70kD protein 9B; peptide-binding protein 74; 75 kDa glucose-regulated protein; catecholamine-regulated protein 40; epididymis secretory sperm binding protein Li
UniProt Protein Name	Stress-70 protein, mitochondrial
UniProt Gene Name	HSPA9
UniProt Synonym Gene Names	GRP75; HSPA9B; mt-HSP70; GRP-75; MOT; PBP74
UniProt Entry Name	GRP75_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/grp75-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222487>