

Rabbit TCEB3C Polyclonal Antibody | anti-TCEB3C antibody

Rabbit anti-TCEB3C Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222575	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TCEB3C; Polyclonal Antibody; Rabbit anti-TCEB3C Antibody; TCEB3L2; RNA polymerase II transcription factor SIII subunit A3; Elongin-A3; EloA3; Transcription elongation factor B polypeptide 3C; anti-TCEB3C antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-TCEB3C antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TCEB3C. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TCEB3C staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of TCEB3C staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TCEB3C expression in Jurkat (A), mouse kidney (B), rat kidney (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-TCEB3C antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	156938279
NCBI GeneID	162699
NCBI Accession #	NP_663628.2
NCBI GenBank Nucleotide #	NM_145653.3
UniProt Accession #	Q8NG57
Molecular Weight	59,744 Da
NCBI Official Full Name	RNA polymerase II transcription factor SIII subunit A3
NCBI Official Synonym Full Names	transcription elongation factor B polypeptide 3C (elongin A3)
NCBI Official Symbol	TCEB3C
NCBI Official Synonym Symbols	HsT829; TCEB3L2
NCBI Protein Information	RNA polymerase II transcription factor SIII subunit A3; eloA3; elongin-A3
UniProt Protein Name	RNA polymerase II transcription factor SIII subunit A3
UniProt Gene Name	TCEB3C
UniProt Synonym Gene Names	TCEB3L2; EloA3
UniProt Entry Name	ELOA3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/tceb3c-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222575>