

Rabbit PLA2G4A Polyclonal Antibody | anti-PLA2G4A antibody

Rabbit anti-PLA2G4A Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222596	Rabbit	Human, Mouse, Monkey	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PLA2G4A; Polyclonal Antibody; Rabbit anti-PLA2G4A Antibody; CPLA2; PLA2G4; Cytosolic phospholipase A2; cPLA2; Phospholipase A2 group IVA; anti-PLA2G4A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-PLA2G4A antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PLA2G4A. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of PLA2G4A staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of PLA2G4A staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PLA2G4A expression in HeLa (A), THP1 (B), NIH3T3 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-PLA2G4A antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	23943920
NCBI GeneID	5321
NCBI Accession #	NP_077734.1
NCBI GenBank Nucleotide #	NM_024420.2
UniProt Accession #	P47712; P47713
Molecular Weight	110kDa
NCBI Official Full Name	cytosolic phospholipase A2
NCBI Official Synonym Full Names	phospholipase A2, group IVA (cytosolic, calcium-dependent)
NCBI Official Symbol	PLA2G4A
NCBI Official Synonym Symbols	PLA2G4; cPLA2-alpha
NCBI Protein Information	cytosolic phospholipase A2; cPLA2; lysophospholipase; phospholipase A2 group IVA; phosphatidylcholine 2-acylhydrolase; calcium-dependent phospholipid-binding protein
UniProt Protein Name	Cytosolic phospholipase A2
UniProt Gene Name	PLA2G4A
UniProt Synonym Gene Names	CPLA2; PLA2G4; cPLA2
UniProt Entry Name	PA24A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/pla2g4a-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222596>