

Rabbit TACSTD2 Polyclonal Antibody | anti-TACSTD2 antibody

Rabbit anti-TACSTD2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222664	Rabbit	Human, Mouse, Rat, Pig	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

TACSTD2; Polyclonal Antibody; Rabbit anti-TACSTD2 Antibody; GA733-1; M1S1; TROP2; Tumor-associated calcium signal transducer 2; Cell surface glycoprotein Trop-2; Membrane component chromosome 1 surface marker 1; Pancreatic carcinoma marker protein GA733-1; anti-TACSTD2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-TACSTD2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TACSTD2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of TACSTD2 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of TACSTD2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of TACSTD2 expression in human prostate (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-TACSTD2 antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	166795236
NCBI GeneID	4070
NCBI Accession #	NP_002344.2
NCBI GenBank Nucleotide #	NM_002353.2
UniProt Accession #	P09758 ; Q8BGV3 ; Q6P9Z6
Molecular Weight	35,709 Da
NCBI Official Full Name	tumor-associated calcium signal transducer 2
NCBI Official Synonym Full Names	tumor-associated calcium signal transducer 2
NCBI Official Symbol	TACSTD2
NCBI Official Synonym Symbols	EGP1; GP50; M1S1; EGP-1; TROP2; GA7331; GA733-1
NCBI Protein Information	tumor-associated calcium signal transducer 2; epithelial glycoprotein-1; cell surface glycoprotein TROP2; cell surface glycoprotein Trop-2; pancreatic carcinoma marker protein GA7331; pancreatic carcinoma marker protein GA733-1; gastrointestinal tumor-ass
UniProt Protein Name	Tumor-associated calcium signal transducer 2
UniProt Gene Name	TACSTD2
UniProt Synonym Gene Names	GA733-1; M1S1; TROP2
UniProt Entry Name	TACD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/tacstd2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222664>