

Rabbit CD222 Polyclonal Antibody | anti-CD222 antibody

Rabbit anti-CD222 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222709	Rabbit	Human, Mouse, Bovine	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD222; Polyclonal Antibody; Rabbit anti-CD222 Antibody; MPRI; Cation-independent mannose-6-phosphate receptor; CI Man-6-P receptor; CI-MPR; M6PR; 300 kDa mannose 6-phosphate receptor; MPR 300; Insulin-like growth factor 2 receptor; Insulin-like growth factor II receptor; IGF-II receptor; M6P/IGF2 receptor; M6P/IGF2R; anti-CD222 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Bovine
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CD222 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human CD222. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CD222 staining in Jurkay cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD222 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD222 expression in Jurkay (A), K562 (B), Caco2 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CD222 antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	119964726
NCBI GeneID	3482
NCBI Accession #	NP_000867.2
NCBI GenBank Nucleotide #	NM_000876.2
UniProt Accession #	P11717 ; Q07113
Molecular Weight	274,375 Da
NCBI Official Full Name	cation-independent mannose-6-phosphate receptor
NCBI Official Synonym Full Names	insulin-like growth factor 2 receptor
NCBI Official Symbol	IGF2R
NCBI Official Synonym Symbols	MPR1; MPRI; CD222; CIMPR; M6P-R
NCBI Protein Information	cation-independent mannose-6-phosphate receptor; M6PR; CI-MPR; MPR 300; M6P/IGF2R; IGF-II receptor; M6P/IGF2 receptor; CI Man-6-P receptor; 300 kDa mannose 6-phosphate receptor; insulin-like growth factor II receptor; cation-independent mannose-6 phosphat
UniProt Protein Name	Cation-independent mannose-6-phosphate receptor
UniProt Gene Name	IGF2R
UniProt Synonym Gene Names	MPRI; CI Man-6-P receptor; CI-MPR; M6PR; MPR 300; IGF-II receptor; M6P/IGF2R
UniProt Entry Name	MPRI_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cd222-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222709>