

Rabbit STAM2 Polyclonal Antibody | anti-STAM2 antibody

Rabbit anti-STAM2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222728	Rabbit	Human, Mouse, Rat	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

STAM2; Polyclonal Antibody; Rabbit anti-STAM2 Antibody; HBP; Signal transducing adapter molecule 2; STAM-2; Hrs-binding protein; anti-STAM2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-STAM2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human STAM2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of STAM2 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of STAM2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of STAM2 expression in HepG2 (A), HeLa (B), NIH3T3 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-STAM2 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	188219625
NCBI GeneID	10254
NCBI Accession #	NP_005834.4
NCBI GenBank Nucleotide #	NM_005843.4
UniProt Accession #	O75886; O88811; Q5XHY7
Molecular Weight	58,164 Da
NCBI Official Full Name	signal transducing adapter molecule 2
NCBI Official Synonym Full Names	signal transducing adaptor molecule (SH3 domain and ITAM motif) 2
NCBI Official Symbol	STAM2
NCBI Official Synonym Symbols	Hbp; STAM2A; STAM2B
NCBI Protein Information	signal transducing adapter molecule 2; STAM-2; HSE1 homolog; Hrs-binding protein; STAM-like protein containing SH3 and ITAM domains 2
UniProt Protein Name	Signal transducing adapter molecule 2
UniProt Gene Name	STAM2
UniProt Synonym Gene Names	HBP; STAM-2
UniProt Entry Name	STAM2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/stam2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222728>