

Rabbit p67 Polyclonal Antibody | anti-p67 antibody

Rabbit anti-p67 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222731	Rabbit	Human, Mouse, Rat, Bovine	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

p67; Polyclonal Antibody; Rabbit anti-p67 Antibody; UNC18A; Syntaxin-binding protein 1; MUNC18-1; N-Sec1; Protein unc-18 homolog 1; Unc18-1; Protein unc-18 homolog A; Unc-18A; anti-p67 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-p67 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human p67. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of p67 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of p67 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of p67 expression in Jurkat (A), HeLa (B), human brain (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-p67 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	73760415
NCBI GeneID	6812
NCBI Accession #	NP_001027392.1
NCBI GenBank Nucleotide #	NM_001032221.3
UniProt Accession #	P61764; O08599; P61765
Molecular Weight	67,569 Da
NCBI Official Full Name	syntaxin-binding protein 1 isoform b
NCBI Official Synonym Full Names	syntaxin binding protein 1
NCBI Official Symbol	STXBP1
NCBI Official Synonym Symbols	P67; NSEC1; UNC18; RBSEC1; MUNC18-1
NCBI Protein Information	syntaxin-binding protein 1; N-Sec1; unc-18A; unc18-1; neuronal SEC1; protein unc-18 homolog 1; protein unc-18 homolog A
UniProt Protein Name	Syntaxin-binding protein 1
UniProt Gene Name	STXBP1
UniProt Synonym Gene Names	UNC18A; Unc18-1; Unc-18A
UniProt Entry Name	STXB1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/p67-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222731>