

Rabbit STEAP2 Polyclonal Antibody | anti-STEAP2 antibody

Rabbit anti-STEAP2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA222776	Rabbit	Human, Mouse, Rat, Bovine	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

STEAP2; Polyclonal Antibody; Rabbit anti-STEAP2 Antibody; PCANAP1; STAMP1; Metalloreductase STEAP2; Prostate cancer-associated protein 1; Protein up-regulated in metastatic prostate cancer; PUMPCn; Six-transmembrane epithelial antigen of prostate 2; SixTransMembrane protein of prostate 1; anti-STEAP2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-STEAP2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human STEAP2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of STEAP2 staining in HEK293T cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of STEAP2 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of STEAP2 expression in HEK293T (A), mouse spleen (B), rat spleen (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-STEAP2 antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	100913194
NCBI GeneID	261729
NCBI Accession #	NP_001035755.1
NCBI GenBank Nucleotide #	NM_001040665.1
UniProt Accession #	Q8NFT2; Q8BWB6
Molecular Weight	56,056 Da
NCBI Official Full Name	metalloreductase STEAP2 isoform a
NCBI Official Synonym Full Names	STEAP family member 2, metalloreductase
NCBI Official Symbol	STEAP2
NCBI Official Synonym Symbols	STMP; IPCA1; PUMPCn; STAMP1; PCANAP1
NCBI Protein Information	metalloreductase STEAP2; prostate cancer associated protein 1; prostate cancer-associated protein 1; SixTransMembrane Protein of Prostate 1; protein upregulated in metastatic prostate cancer; protein up-regulated in metastatic prostate cancer; six transme
UniProt Protein Name	Metalloreductase STEAP2
UniProt Gene Name	STEAP2
UniProt Synonym Gene Names	PCANAP1; STAMP1; PUMPCn
UniProt Entry Name	STEAP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/steap2-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/222776>