

Rabbit PDGFR alpha Polyclonal Antibody | anti-PDGFRalpha antibody

Rabbit anti-PDGFR alpha Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223282	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Dog, Monkey, Pig	Immunofluorescence, Western Blot


PRODUCT SYNONYMS

PDGFR alpha; Polyclonal Antibody; Rabbit anti-PDGFR alpha Antibody; PDGFR2; RHEPDGFRA; Platelet-derived growth factor receptor alpha; PDGF-R-alpha; PDGFR-alpha; Alpha platelet-derived growth factor receptor; Alpha-type platelet-derived growth factor receptor; CD140 antigen-like family member A; CD140a antigen; Platelet-derived growth factor alpha receptor; Platelet-derived growth factor receptor 2; PDGFR-2; CD140a; anti-PDGFRalpha antibody


PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Dog, Monkey, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-PDGFRalpha antibody	IC (Immunochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PDGFR alpha. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES


IF (Immunofluorescence)

Immunofluorescent analysis of PDGFR alpha staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).


IHC (Immunohistochemistry)



WB (Western Blot)

Western blot analysis of PDGFR alpha expression in HepG2 (A), NIH3T3 (B), Saos2 (C), HeLa (D), A431 (E) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-PDGFRalpha antibody	Primary antibodies
--	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	5453870
NCBI GeneID	5156
NCBI Accession #	NP_006197.1
NCBI GenBank Nucleotide #	NM_006206.4
UniProt Accession #	P16234; P26618; P20786
Molecular Weight	122,670 Da
NCBI Official Full Name	platelet-derived growth factor receptor alpha
NCBI Official Synonym Full Names	platelet-derived growth factor receptor, alpha polypeptide
NCBI Official Symbol	PDGFRA
NCBI Official Synonym Symbols	CD140A; PDGFR2; PDGFR-2; RHEPDGFRA
NCBI Protein Information	platelet-derived growth factor receptor alpha; PDGF-R-alpha; CD140a antigen; PDGFRA/BCR fusion; CD140 antigen-like family member A; platelet-derived growth factor receptor 2; alpha-type platelet-derived growth factor receptor; rearranged-in-hypereosinophi
UniProt Protein Name	Platelet-derived growth factor receptor alpha
UniProt Gene Name	PDGFRA
UniProt Synonym Gene Names	PDGFR2; RHEPDGFRA; PDGF-R-alpha; PDGFR-alpha; PDGFR-2
UniProt Entry Name	PGFRA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/pdgr-alpha-antibody-immunochemistry-immunofluorescence-western-blot-elisa/223282>