

Rabbit c-Met Polyclonal Antibody | anti-c-Met antibody

Rabbit anti-c-Met Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223529	Rabbit	Human, Mouse, Rat, Rabbit, Sheep	Immunofluorescence, Western Blot


PRODUCT SYNONYMS

c-Met; Polyclonal Antibody; Rabbit anti-c-Met Antibody; Hepatocyte growth factor receptor; HGF receptor; HGF/SF receptor; Proto-oncogene c-Met; Scatter factor receptor; SF receptor; Tyrosine-protein kinase Met; anti-c-Met antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Rabbit, Sheep
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-c-Met antibody	IC (Immunocytochemistry), IF (Immunofluorescence), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human c-Met. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of c-Met staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



WB (Western Blot)

Western blot analysis of c-Met expression in HeLa (A), A431 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-c-Met antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	42741655
NCBI GeneID	4233
NCBI Accession #	NP_000236.2
NCBI GenBank Nucleotide #	NM_000245.2
UniProt Accession #	P08581; P16056; P97523
Molecular Weight	155,541 Da
NCBI Official Full Name	hepatocyte growth factor receptor isoform b
NCBI Official Synonym Full Names	met proto-oncogene (hepatocyte growth factor receptor)
NCBI Official Symbol	MET
NCBI Official Synonym Symbols	HGFR; AUTS9; RCCP2; c-Met
NCBI Protein Information	hepatocyte growth factor receptor; SF receptor; HGF receptor; HGF/SF receptor; proto-oncogene c-Met; scatter factor receptor; tyrosine-protein kinase Met; met proto-oncogene tyrosine kinase
UniProt Protein Name	Hepatocyte growth factor receptor
UniProt Gene Name	MET
UniProt Synonym Gene Names	HGF receptor; SF receptor
UniProt Entry Name	MET_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/c-met-antibody-immunochemistry-immunofluorescence-western-blot-elisa/223529>