

Rabbit EGR Polyclonal Antibody | anti-EGR antibody

Rabbit anti-EGR Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223671	Rabbit	Human, Mouse, Rat, Bovine, Chicken	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

EGR; Polyclonal Antibody; Rabbit anti-EGR Antibody; EGR1; KROX24; ZNF225; Early growth response protein 1; EGR-1; AT225; Nerve growth factor-induced protein A; NGFI-A; Transcription factor ETR103; Transcription factor Zif268; Zinc finger protein 225; Zinc finger protein Krox-24; EGR2; KROX20; E3 SUMO-protein ligase EGR2; AT591; Early growth response protein 2; EGR-2; Zinc finger protein Krox-20; anti-EGR antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-EGR antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human EGR1. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of EGR staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of EGR staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of EGR expression in MCF7 (A), SHSY5Y (B), PC12 (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-EGR antibody [Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4503493
NCBI GeneID	1958
NCBI Accession #	NP_001955.1
NCBI GenBank Nucleotide #	NM_001964.2
UniProt Accession #	P18146 ; P11161 ; P08046 ; P08152 ; P08154 ; P51774
Molecular Weight	57,507 Da
NCBI Official Full Name	early growth response protein 1
NCBI Official Synonym Full Names	early growth response 1
NCBI Official Symbol	EGR1
NCBI Official Synonym Symbols	TIS8; AT225; G0S30; NGFI-A; ZNF225; KROX-24; ZIF-268
NCBI Protein Information	early growth response protein 1; EGR-1; zinc finger protein 225; transcription factor ETR103; transcription factor Zif268; zinc finger protein Krox-24; nerve growth factor-induced protein A
UniProt Protein Name	Early growth response protein 1
UniProt Gene Name	EGR1
UniProt Synonym Gene Names	KROX24; ZNF225; EGR-1; NGFI-A
UniProt Entry Name	EGR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/egr-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223671>