

Rabbit HADH2 Polyclonal Antibody | anti-HADH2 antibody

Rabbit anti-HADH2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223687	Rabbit	Human, Mouse, Rat, Bovine, Dog, Pig	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

HADH2; Polyclonal Antibody; Rabbit anti-HADH2 Antibody; ERAB; MRPP2; SCHAD; XH98G2; 3-hydroxyacyl-CoA dehydrogenase type-2; 17-beta-hydroxysteroid dehydrogenase 10; 17-beta-HSD 10; 3-hydroxy-2-methylbutyryl-CoA dehydrogenase; 3-hydroxyacyl-CoA dehydrogenase type II; Endoplasmic reticulum-associated amyloid beta-peptide-binding protein; Mitochondrial ribonuclease P protein 2; Mitochondrial RNase P protein 2; Short-chain type dehydrogenase/reductase XH98G2; Type II HADH; anti-HADH2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-HADH2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HADH2. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of HADH2 staining in SKNSH cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of HADH2 staining in human lung formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)
Western blot analysis of HADH2 expression in SKNSH (A), HEK293T (B), rat brain (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-HADH2 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	83715985
NCBI GeneID	3028
NCBI Accession #	NP_001032900.1
NCBI GenBank Nucleotide #	NM_001037811.2
UniProt Accession #	Q99714; O08756; O70351
Molecular Weight	26,923 Da
NCBI Official Full Name	3-hydroxyacyl-CoA dehydrogenase type-2 isoform 2
NCBI Official Synonym Full Names	hydroxysteroid (17-beta) dehydrogenase 10
NCBI Official Symbol	HSD17B10
NCBI Official Synonym Symbols	ABAD; CAMR; ERAB; HCD2; MHBD; HADH2; MRPP2; MRX17; MRX31; SCHAD; MRXS10; SDR5C1; 17b-HSD10; DUPXp11.22
NCBI Protein Information	3-hydroxyacyl-CoA dehydrogenase type-2; mitochondrial RNase P subunit 2; AB-binding alcohol dehydrogenase; mitochondrial ribonuclease P protein 2; 3-hydroxy-2-methylbutyryl-CoA dehydrogenase; short chain type dehydrogenase/reductase XH98G2; amyloid-beta p
UniProt Protein Name	3-hydroxyacyl-CoA dehydrogenase type-2
UniProt Gene Name	HSD17B10
UniProt Synonym Gene Names	ERAB; HADH2; MRPP2; SCHAD; XH98G2; 17-beta-HSD 10; Mitochondrial RNase P protein 2
UniProt Entry Name	HCD2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.