

Rabbit c-SRC Polyclonal Antibody | anti-c-SRC antibody

Rabbit anti-c-SRC Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223753	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Pig, Zebrafish	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

c-SRC; Polyclonal Antibody; Rabbit anti-c-SRC Antibody; SRC1; Proto-oncogene tyrosine-protein kinase Src; Proto-oncogene c-Src; pp60c-src; p60-Src; anti-c-SRC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Pig, Zebrafish
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-c-SRC antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human c-SRC. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of c-SRC staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of c-SRC staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of c-SRC expression in HeLa (A), HuvEc (B), Jurkat (C), COLO205 (D) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-c-SRC antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885609
NCBI GeneID	6714
NCBI Accession #	NP_005408.1
NCBI GenBank Nucleotide #	NM_005417.3
UniProt Accession #	P12931 ; P05480 ; Q9WUD9
Molecular Weight	59,835 Da
NCBI Official Full Name	proto-oncogene tyrosine-protein kinase Src
NCBI Official Synonym Full Names	v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog (avian)
NCBI Official Symbol	SRC
NCBI Official Synonym Symbols	ASV; SRC1; c-SRC; p60-Src
NCBI Protein Information	proto-oncogene tyrosine-protein kinase Src; proto-oncogene c-Src; tyrosine kinase pp60c-src; tyrosine-protein kinase SRC-1; protooncogene SRC, Rous sarcoma
UniProt Protein Name	Proto-oncogene tyrosine-protein kinase Src
UniProt Gene Name	SRC
UniProt Synonym Gene Names	SRC1; p60-Src
UniProt Entry Name	SRC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/c-src-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223753>