

Rabbit p63 Polyclonal Antibody | anti-p63 antibody

Rabbit anti-p63 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223772	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Dog, Monkey, Pig	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

p63; Polyclonal Antibody; Rabbit anti-p63 Antibody; KET; P63; P73H; P73L; TP73L; Tumor protein 63; Chronic ulcerative stomatitis protein; CUSP; Keratinocyte transcription factor KET; Transformation-related protein 63; TP63; Tumor protein p73-like; p73L; p40; p51; anti-p63 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Dog, Monkey, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-p63 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human p63. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of p63 staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of p63 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of p63 expression in A549 (A), HCT116 (B), COLO205 (C), LOVO (D), HT29 (E) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-p63 antibody	Primary antibodies
---	--------------------

NCBI AND UNIPROT INFORMATION

NCBI GI #	169234657
NCBI GeneID	8626
NCBI Accession #	NP_001108450.1
NCBI GenBank Nucleotide #	NM_001114978.1
UniProt Accession #	Q9H3D4; O88898; Q9JJP6
Molecular Weight	76,785 Da
NCBI Official Full Name	tumor protein 63 isoform 2
NCBI Official Synonym Full Names	tumor protein p63
NCBI Official Symbol	TP63
NCBI Official Synonym Symbols	AIS; KET; LMS; NBP; RHS; p40; p51; p63; EEC3; OFC8; p73H; p73L; SHFM4; TP53L; TP73L; p53CP; TP53CP; B(p51A); B(p51B)
NCBI Protein Information	tumor protein 63; CUSP; transformation-related protein 63; tumor protein p53-competing protein; amplified in squamous cell carcinoma; chronic ulcerative stomatitis protein; keratinocyte transcription factor KET; tumor protein p63 deltaN isoform delta
UniProt Protein Name	Tumor protein 63
UniProt Gene Name	TP63
UniProt Synonym Gene Names	KET; P63; P73H; P73L; TP73L; p63; CUSP; TP63; p73L
UniProt Entry Name	P63_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
 Online datasheet: <https://www.aaabiotech.com/polyclonal/p63-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223772>