

Rabbit CD256 Polyclonal Antibody | anti-CD256 antibody

Rabbit anti-CD256 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223773	Rabbit	Human, Mouse, Rat, Bovine, Dog, Pig	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

CD256; Polyclonal Antibody; Rabbit anti-CD256 Antibody; APRIL; TALL2; ZTNF2; Tumor necrosis factor ligand superfamily member 13; A proliferation-inducing ligand; TNF- and APOL-related leukocyte expressed ligand 2; TALL-2; TNF-related death ligand 1; TRDL-1; anti-CD256 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Pig
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-CD256 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD256. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of CD256 staining in H9C2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of CD256 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of CD256 expression in HEK293T (A), Hela (B), H446 (C), mouse brain (D), mouse kidney (E), rat brain (F), rat kidney (G) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-CD256 antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	310750387
NCBI GeneID	8741
NCBI Accession #	NP_001185552.1
NCBI GenBank Nucleotide #	NM_001198623.1
UniProt Accession #	O75888; Q9D777
Molecular Weight	27,433 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 13 isoform zeta
NCBI Official Synonym Full Names	tumor necrosis factor (ligand) superfamily, member 13
NCBI Official Symbol	TNFSF13
NCBI Official Synonym Symbols	APRIL; CD256; TALL2; ZTNF2; TALL-2; TRDL-1; UNQ383/PRO715
NCBI Protein Information	tumor necrosis factor ligand superfamily member 13; a proliferation-inducing ligand; tumor necrosis factor-like protein ZTNF2; tumor necrosis factor-related death ligand-1; TNF- and APOL-related leukocyte expressed ligand 2
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 13
UniProt Gene Name	TNFSF13
UniProt Synonym Gene Names	APRIL; TALL2; ZTNF2; APRIL; TALL-2; TRDL-1
UniProt Entry Name	TNF13_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/cd256-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223773>