

Rabbit ERK3 Polyclonal Antibody | anti-ERK3 antibody

Rabbit anti-ERK3 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223827	Rabbit	Human, Mouse, Bovine, Chicken, Monkey	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

ERK3; Polyclonal Antibody; Rabbit anti-ERK3 Antibody; PRKM6; Mitogen-activated protein kinase 6; MAP kinase 6; MAPK 6; Extracellular signal-regulated kinase 3; ERK-3; MAP kinase isoform p97; p97-MAPK; anti-ERK3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Bovine, Chicken, Monkey
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-ERK3 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ERK3. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of ERK3 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of ERK3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of ERK3 expression in HeLa (A), NIH3T3 (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-ERK3 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506091
NCBI GeneID	5597
NCBI Accession #	NP_002739.1
NCBI GenBank Nucleotide #	NM_002748.3
UniProt Accession #	Q16659; Q61532
Molecular Weight	82,681 Da
NCBI Official Full Name	mitogen-activated protein kinase 6
NCBI Official Synonym Full Names	mitogen-activated protein kinase 6
NCBI Official Symbol	MAPK6
NCBI Official Synonym Symbols	ERK3; PRKM6; p97MAPK; HsT17250
NCBI Protein Information	mitogen-activated protein kinase 6; ERK-3; MAPK 6; p97-MAPK; MAP kinase 6; MAP kinase isoform p97; protein kinase, mitogen-activated 5; protein kinase, mitogen-activated 6; extracellular signal-regulated kinase 3; extracellular signal-regulated kinase, p9
UniProt Protein Name	Mitogen-activated protein kinase 6
UniProt Gene Name	MAPK6
UniProt Synonym Gene Names	ERK3; PRKM6; MAP kinase 6; MAPK 6; ERK-3; p97-MAPK
UniProt Entry Name	MK06_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/erk3-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223827>