

Rabbit GAP43 Polyclonal Antibody | anti-GAP43 antibody

Rabbit anti-GAP43 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223877	Rabbit	Human, Mouse, Rat, Bovine, Chicken, Monkey, Zebrafish	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

GAP43; Polyclonal Antibody; Rabbit anti-GAP43 Antibody; Neuromodulin; Axonal membrane protein GAP-43; Growth-associated protein 43; Neural phosphoprotein B-50; pp46; anti-GAP43 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Chicken, Monkey, Zebrafish
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-GAP43 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GAP43. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of GAP43 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of GAP43 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of GAP43 expression in HeLa (A), mouse brain (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-GAP43 antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	194248057
NCBI GeneID	2596
NCBI Accession #	NP_001123536.1
NCBI GenBank Nucleotide #	NM_001130064.1
UniProt Accession #	P17677; P06837; P07936
Molecular Weight	24,803 Da
NCBI Official Full Name	neuromodulin isoform 1
NCBI Official Synonym Full Names	growth associated protein 43
NCBI Official Symbol	GAP43
NCBI Official Synonym Symbols	B-50; PP46
NCBI Protein Information	neuromodulin; protein F1; neural phosphoprotein B-50; growth-associated protein 43; axonal membrane protein GAP-43; calmodulin-binding protein P-57; nerve growth-related peptide GAP43; neuron growth-associated protein 43
UniProt Protein Name	Neuromodulin
UniProt Gene Name	GAP43
UniProt Entry Name	NEUM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/gap43-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223877>