

Rabbit PPAR gamma Polyclonal Antibody | anti-PPARgamma antibody

Rabbit anti-PPAR gamma Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA223965	Rabbit	Human, Mouse, Rat, Dog, Rabbit	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

PPAR gamma; Polyclonal Antibody; Rabbit anti-PPAR gamma Antibody; NR1C3; Peroxisome proliferator-activated receptor gamma; PPAR-gamma; Nuclear receptor subfamily 1 group C member 3; anti-PPARgamma antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Dog, Rabbit
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-PPARgamma antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PPAR gamma. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of PPAR gamma staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.



IHC (Immunohistochemistry)

Immunohistochemical analysis of PPAR gamma staining in human colon formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of PPAR gamma expression in HeLa (A), MCF7 (B), mouse brain (C) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-PPARgamma antibody

[Primary antibodies](#)

NCBI AND UNIPROT INFORMATION

NCBI GI #	116284368
NCBI GeneID	5468
NCBI Accession #	NP_005028.4
NCBI GenBank Nucleotide #	NM_005037.5
UniProt Accession #	P37231 ; P37238 ; O88275
Molecular Weight	57,620 Da
NCBI Official Full Name	peroxisome proliferator-activated receptor gamma isoform 1
NCBI Official Synonym Full Names	peroxisome proliferator-activated receptor gamma
NCBI Official Symbol	PPARG
NCBI Official Synonym Symbols	GLM1; CIMT1; NR1C3; PPARG1; PPARG2; PPARGgamma
NCBI Protein Information	peroxisome proliferator-activated receptor gamma; PPAR gamma; PPAR-gamma; nuclear receptor subfamily 1 group C member 3; peroxisome proliferator-activated receptor gamma 1; peroxisome proliferator-activated nuclear receptor gamma variant 1
UniProt Protein Name	Peroxisome proliferator-activated receptor gamma
UniProt Gene Name	PPARG
UniProt Synonym Gene Names	NR1C3; PPAR-gamma
UniProt Entry Name	PPARG_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/ppar-gamma-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/223965>