

Rabbit eNOS Polyclonal Antibody | anti-eNOS antibody

Rabbit anti-eNOS Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA224022	Rabbit	Human, Mouse, Rat, Bovine, Dog, Pig, Rabbit, Sheep	Immunofluorescence, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

eNOS; Polyclonal Antibody; Rabbit anti-eNOS Antibody; Nitric oxide synthase; endothelial; Constitutive NOS; cNOS; EC-NOS; Endothelial NOS; NOS type III; NOSIII; anti-eNOS antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat, Bovine, Dog, Pig, Rabbit, Sheep
Clonality	Polyclonal
Purity/Purification	The antibody was purified by immunogen affinity chromatography.
Form/Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Concentration	1mg/mL (varies by lot)
Applicable Applications for anti-eNOS antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human eNOS. The exact sequence is proprietary.
Preparation and Storage	Store at -20 degree C. Avoid repeated freeze/thaw cycles.

IMAGES




IF (Immunofluorescence)

Immunofluorescent analysis of eNOS staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



IHC (Immunohistochemistry)

Immunohistochemical analysis of eNOS staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



WB (Western Blot)

Western blot analysis of eNOS expression in HepG2 (A), Hela (B) whole cell lysates.

ADDITIONAL INFORMATION

Product Categories/Family for anti-eNOS antibody

Primary antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	40254422
NCBI GeneID	4846
NCBI Accession #	NP_000594.2
NCBI GenBank Nucleotide #	NM_000603.4
UniProt Accession #	P29474; P70313; Q62600
Molecular Weight	133,289 Da
NCBI Official Full Name	nitric oxide synthase, endothelial isoform 1
NCBI Official Synonym Full Names	nitric oxide synthase 3 (endothelial cell)
NCBI Official Symbol	NOS3
NCBI Official Synonym Symbols	eNOS; ECNOS
NCBI Protein Information	nitric oxide synthase, endothelial; cNOS; EC-NOS; NOSIII; NOS type III; endothelial NOS; constitutive NOS
UniProt Protein Name	Nitric oxide synthase, endothelial
UniProt Gene Name	NOS3
UniProt Synonym Gene Names	cNOS; eNOS; NOSIII
UniProt Entry Name	NOS3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/enos-antibody-immunofluorescence-immunohistochemistry-western-blot-elisa/224022>