

S-acetyl-PEG5-alcohol Biochemical

S-acetyl-PEG5-alcohol

CATALOG #
AAA227405


PRODUCT SYNONYMS

S-acetyl-PEG5-alcohol; S-acetyl-PEG5-alcohol biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	1000415-61-1
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
In vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins[1].
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure
CefaloniumM/hydrate, CAS 1486466-27-6



Structure
CefaloniumM/hydrate, CAS 1486466-27-6

ADDITIONAL INFORMATION

Related Product Information for S-acetyl-PEG5-alcohol biochemical	S-acetyl-PEG5-alcohol is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.
References	1. Harada K, et, al. Assessment of the Usefulness of Cefapirin and Cefalonium Disks for Susceptibility Testing of Staphylococcus aureus Isolates from Bovine Mastitis. Antibiotics (Basel). 2020 Apr 21;9(4):197. 2. Demon D, et, al. The intramammary efficacy of first generation cephalosporins against Staphylococcus aureus mastitis in mice. Vet Microbiol. 2012 Nov 9;160(1-2):141-50. 3. Berry EA, et, al. Effect of an intramammary teat seal and dry cow antibiotic in relation to dry period length on postpartum mastitis. J Dairy Sci. 2007 Feb;90(2):760-5.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/biochemical/s-acetyl-peg5-alcohol/227405>