

m-PEG9-Br biochemical

m-PEG9-Br

CATALOG #
AAA227546


PRODUCT SYNONYMS

m-PEG9-Br; CCK Octapeptide; desulfated TFA; m-PEG9-Br biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	125562-30-3
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
In vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins[1]
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES


Structure

Cholecystokinin Octapeptide, desulfated TFA, CAS 171486-94-5


Structure

Cholecystokinin Octapeptide, desulfated TFA, CAS 171486-94-5

ADDITIONAL INFORMATION

Related Product Information for m-PEG9-Br biochemical	m-PEG9-Br is a PEG-based PROTAC linker that can be used in the synthesis of PROTACs.
References	1. L R Johnson, et al. Effect of sulfation on the gastrointestinal actions of caerulein. Gastroenterology. 1970 Feb;58(2):208-16.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.