

CYH33 methanesulfonate Biochemical

CYH33 methanesulfonate

CATALOG #
AAA227756


PRODUCT SYNONYMS

CYH33 methanesulfonate; CYH33 methanesulfonate biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	1494684-33-1
Targets&IC50	PI3Kgamma:225 nM (IC50), PI3Kalpha:5.9 nM (IC50), PI3Kbeta:598 nM (IC50), PI3Kdelta:78.7 nM (IC50)
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
In vitro	CYH33 (2-20 mg/kg; oral; once a day for 21 days) methanesulfonate potently restrains tumor growth in mice bearing human breast cancer cell xenografts[2]. Single administration of CYH33 (20mg/kg; oral) methanesulfonate significantly down-regulates the level of phosphorylated Akt in tumor tissues, demonstrating the suppression of PI3K signaling in nude mice[2]. CYH33 (10mg/kg; oral; once a day for 18-d or 20-d respectively) methanesulfonate delays the restoration of blood glucose and area under the curve (AUC) of blood glucose increased upon CYH33 treatment in T47D xenografts and R26-Pik3ca H1047R ;MMTV-Cre mice[2]. Animal Model: SCID mice aged 4-6 weeks bearing human breast cancer T47D xenografts[2]. Dosage: 2, 5, 10, 20 mg/kg. Administration: Oral; once a day for 21 days. Result: Displayed marginal inhibitory effect on the tumor growth at lower doses (2 and 5mg/kg) and significantly attenuated tumor growth at the dose of 10 or 20mg/kg, yielding T/C values of 58.36% and 49.42% respectively.
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure
PYBG-TMR, CAS 2067339-51-7



Structure
PYBG-TMR, CAS 2067339-51-7

ADDITIONAL INFORMATION

Related Product Information for CYH33 methanesulfonate biochemical	CYH33 methanesulfonate is an orally active, highly selective PI3Kalpha inhibitor with IC 50 s of 5.9 nM/598 nM/78.7 nM/225 nM against alpha/beta/delta/gamma isoform, respectively. CYH33 methanesulfonate inhibits phosphorylation of Akt, ERK and induces significant G1 phase arrest in breast cancer cells and non-small cell lung cancer (NSCLC) cells. CYH33 methanesulfonate has potent activity against solid tumors.
References	1. Xinbo Song, et al. Terminal alkyne substituted O 6 -benzylguanine for versatile and effective syntheses of fluorescent labels to genetically encoded SNAP-tags. RSC Advances. 26 Feb 2015.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/biochemical/cyh33-methanesulfonate/227756>