

(S,R,S)-AHPC-C8-NH2 hydrochloride Biochemical

(S,R,S)-AHPC-C8-NH2 hydrochloride

CATALOG #
AAA228363


PRODUCT SYNONYMS

(S; R; S)-AHPC-C8-NH2 hydrochloride; S)-AHPC-C8-NH2 hydrochloride biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	2376139-49-8
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
In vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins[2].
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure
BMS-986121, CAS 313671-26-0



Structure
BMS-986121, CAS 313671-26-0

ADDITIONAL INFORMATION

Related Product Information for (S,R,S)-AHPC-C8-NH2 hydrochloride biochemical	(S,R,S)-AHPC-C8-NH2 (VH032-C8-NH2) hydrochloride is a synthesized E3 ligase ligand-linker conjugate that incorporates the VH032 based VHL ligand and a linker used in PROTAC technology.
References	- Burford NT, et al. Discovery of positive allosteric modulators and silent allosteric modulators of the u-opioid receptor. Proc Natl Acad Sci U S A. 2013;110(26):10830-10835. - WO2014107344 - Bisignano P, et al. Ligand-Based Discovery of a New Scaffold for Allosteric Modulation of the u-Opioid Receptor. J Chem Inf Model. 2015;55(9):1836-1843.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/biochemical/srs-ahpc-c8-nh2-hydrochloride/228363>