

OICR-9429-N-C2-NH2 Biochemical

OICR-9429-N-C2-NH2

CATALOG #
AAA228397


PRODUCT SYNONYMS

OICR-9429-N-C2-NH2; OICR-9429-N-C2-NH2 biochemical


PRODUCT SPECIFICATIONS

Purity/Purification	98%
CAS	2407457-55-8
Solubility	<1mg/ml refers to the product slightly soluble or insoluble
In vitro	PROTACs contain two different ligands connected by a linker; one is a ligand for an E3 ubiquitin ligase and the other is for the target protein. PROTACs exploit the intracellular ubiquitin-proteasome system to selectively degrade target proteins[2].
Preparation and Storage	Powder: -20 degree C for 3 years In solvent: -80 degree C for 2 years

IMAGES



Structure
PEG-8 laurate, CAS 35179-86-3



Structure
PEG-8 laurate, CAS 35179-86-3

ADDITIONAL INFORMATION

Related Product Information for OICR-9429-N-C2-NH2 biochemical	OICR-9429-N-C2-NH2 is a ligand for Wd40 repeat domain protein 5 (WDR5) extracted from patent WO2019246570A1, intermediate 2. OICR-9429-N-C2-NH2 can be used in the synthesis of PROTACs.
References	- Bergh BA, et, al. Elasticity of vesicles affects hairless mouse skin structure and permeability. J Control Release. 1999 Dec 6;62(3):367-79. - Barbosa RM, et, al. Physicochemical characterization of surfactant incorporating vesicles that incorporate colloidal magnetite. J Liposome Res. 2013 Mar;23(1):47-53.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

