

Rabbit anti-Human ATG7 Polyclonal Antibody | anti-ATG7 antibody

ATG7 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA229104	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

ATG7; Polyclonal Antibody; ATG7 Antibody; Ubiquitin-like modifier-activating enzyme ATG7; ATG12-activating enzyme E1 ATG7; Autophagy-related protein 7; APG7-like; hAGP7; Ubiquitin-activating enzyme E1-like protein; APG7L; anti-ATG7 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%,Protein G purified
Form/Format	Liquid
Sequence Length	676
Applicable Applications for anti-ATG7 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Ubiquitin-like modifier-activating enzyme ATG7 protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-8668 / sc-12497 / sc-33211
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES




WB (Western Blot)

Western BlotPositive WB detected in: HEK293 whole cell lysateAll lanes: ATG7 antibody at 3.5ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 78,76,69 KDaObserved band size: 78 KDa



IF (Immunofluorescence)

Immunofluorescent analysis of U251 cells using AAA229104 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human cervical cancer using AAA229104 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-ATG7 antibody

E1-like activating enzyme involved in the 2 ubiquitin-like systems required for cytoplasm to vacuole transport (Cvt) and autophagy. Activates ATG12 for its conjugation with ATG5 as well as the ATG8 family proteins for their conjugation with phosphatidylethanolamine. Both systems are needed for the ATG8 association to Cvt vesicles and autophagosomes membranes. Required for autophagic death induced by caspase-8 inhibition. Required for mitophagy which contributes to regulate mitochondrial quantity and quality by eliminating the mitochondria to a basal level to fulfill cellular energy requirements and preventing excess ROS production. Modulates p53/TP53 activity to regulate cell cycle and survival during metabolic stress. Plays also a key role in the maintenance of axonal homeostasis, the prevention of axonal degeneration, the maintenance of hematopoietic stem cells, the formation of Paneth cell granules, as well as in adipose differentiation.

NCBI AND UNIPROT INFORMATION

NCBI GI #	222144227
NCBI GeneID	10533
NCBI Accession #	NP_001129503.2
NCBI GenBank Nucleotide #	NM_001136031.2
UniProt Accession #	O95352; Q7L8L0; Q9BWP2; Q9UFH4; B4E170; E9PB95
Molecular Weight	68,618 Da
NCBI Official Full Name	ubiquitin-like modifier-activating enzyme ATG7 isoform b
NCBI Official Synonym Full Names	autophagy related 7
NCBI Official Symbol	ATG7
NCBI Official Synonym Symbols	GSA7; APG7L; APG7-LIKE
NCBI Protein Information	ubiquitin-like modifier-activating enzyme ATG7
UniProt Protein Name	Ubiquitin-like modifier-activating enzyme ATG7
UniProt Gene Name	ATG7
UniProt Synonym Gene Names	APG7L; APG7-like; hAGP7
UniProt Entry Name	ATG7_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/atg7-antibody-immunofluorescence-immunohistochemistry-elisa/229104>