

Rabbit anti-Human ADM Polyclonal Antibody | anti-ADM antibody

Rabbit anti-human ADM polyclonal Antibody

CATALOG # AAA229111	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

ADM; Polyclonal Antibody; Rabbit anti-human ADM polyclonal Antibody; ProAM N-terminal 20 peptide; PAMP; ProAM-N20; anti-ADM antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	185
Applicable Applications for anti-ADM antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human ADM protein(45-92aa)
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-16495 / sc-16496 / sc-33187
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human kidney tissue using AAA229111 at dilution of 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human lung tissue using AAA229111 at dilution 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-ADM antibody	AM and PAMP are potent hypotensive and vasodilator agents. Numerous actions have been reported most related to the physiologic control of fluid and electrolyte homeostasis. In the kidney, am is diuretic and natriuretic, and both am and pamp inhibit aldosterone secretion by direct adrenal actions. In pituitary gland, both peptides at physiologically relevant doses inhibit basal ACTH secretion. Both peptides appear to act in brain and pituitary gland to facilitate the loss of plasma volume, actions which complement their hypotensive effects in blood vessels. Adrenomedullin, a hypotensive peptide found in human pheochromocytoma, consists of 52 amino acids, has 1 intramolecular disulfide bond, and shows a slight homology with the calcitonin gene-related peptide. It may function as a hormone in circulation control because it is found in blood in a considerable concentration. The precursor, called preproadrenomedullin, is 185 amino acids long. By RNA-blot analysis, human adrenomedullin mRNA was found to be highly expressed in several tissues. Genomic ADM DNA consists of 4 exons and 3 introns, with the 5-prime flanking region containing TATA, CAAT, and GC boxes. There are also multiple binding sites for activator protein-2 and a cAMP-regulated enhancer element.
Product Categories/Family for anti-ADM antibody	Neuroscience

NCBI AND UNIPROT INFORMATION

NCBI GI #	4501945
NCBI GeneID	133
NCBI Accession #	NP_001115.1
NCBI GenBank Nucleotide #	NM_001124.2
UniProt Accession #	P35318; Q6FGW2; B2R793; D3DQV3
Molecular Weight	20,420 Da
NCBI Official Full Name	ADM
NCBI Official Synonym Full Names	adrenomedullin
NCBI Official Symbol	ADM
NCBI Official Synonym Symbols	AM; PAMP
NCBI Protein Information	ADM
UniProt Protein Name	ADM
UniProt Gene Name	ADM
UniProt Synonym Gene Names	AM; AM; PAMP; ProAM-N20
UniProt Entry Name	ADML_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/adm-antibody-immunohistochemistry-elisa/229111>