

Rabbit anti-Human TLR6 Polyclonal Antibody | anti-TLR6 antibody

TLR6 Antibody

CATALOG # AAA229260	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TLR6; Polyclonal Antibody; TLR6 Antibody; Toll-like receptor 6; CD_antigen: CD286; anti-TLR6 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Affinity Purified
Form/Format	Liquid
Sequence Length	796
Applicable Applications for anti-TLR6 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant human Toll-like receptor 6 protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-5658 / sc-5657 / sc-30001
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human thyroid tissue using AAA229260 at dilution of 1:100



WB (Western Blot)

Western blotAll lanes: TLR6 antibody at 5ug/ml+ Jurkat whole cell lysateGoat polyclonal to rabbit at 1/10000 dilutionPredicted band size: 92,56 kDaObserved band size: 92 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-TLR6 antibody	Participates in the innate immune response to bacteria and fungi. Specifically recognizes diacylated and, to a lesser extent, triacylated lipopeptides. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Recognizes mycoplasmal macrophage-activating lipopeptide-2kD (MALP-2), soluble tuberculosis factor (STF), phenol-soluble modulin (PSM) and B.burgdorferi outer surface protein A lipoprotein (OspA-L) cooperatively with TLR2 (PubMed:11441107). In complex with TLR4, promotes sterile inflammation in monocytes/macrophages in response to oxidized low-density lipoprotein (oxLDL) or amyloid-beta 42. In this context, the initial signal is provided by oxLDL- or amyloid-beta 42-binding to CD36. This event induces the formation of a heterodimer of TLR4 and TLR6, which is rapidly internalized and triggers inflammatory response, leading to the NF-kappa-B-dependent production of CXCL1, CXCL2 and CCL9 cytokines, via MYD88 signaling pathway, and CCL5 cytokine, via TICAM1 signaling pathway, as well as IL1B secretion.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	20143971
NCBI GeneID	10333
NCBI Accession #	NP_006059.2
NCBI GenBank Nucleotide #	NM_006068.4
UniProt Accession #	Q9Y2C9; Q2NKL3; B3Y640; B6CH35; B6RFS4; B6RFS5
Molecular Weight	55,828 Da
NCBI Official Full Name	toll-like receptor 6
NCBI Official Synonym Full Names	toll like receptor 6
NCBI Official Symbol	TLR6
NCBI Official Synonym Symbols	CD286
NCBI Protein Information	toll-like receptor 6
UniProt Protein Name	Toll-like receptor 6
UniProt Gene Name	TLR6
UniProt Entry Name	TLR6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/tlr6-antibody-immunohistochemistry-western-blot-elisa/229260>