

Rabbit anti-Human Parathyroid hormone Polyclonal Antibody | anti-PTH antibody

Rabbit anti-human Parathyroid hormone polyclonal Antibody

CATALOG # AAA229512	HOST Rabbit	REACTIVITY Human	APPLICATIONS Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

Parathyroid hormone; Polyclonal Antibody; Rabbit anti-human Parathyroid hormone polyclonal Antibody; Parathyroid hormone protein; Parathormone; Parathyrin; PTH; anti-PTH antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Form/Format	Liquid
Sequence Length	115
Applicable Applications for anti-PTH antibody	WB (Western Blot), ELISA
Immunogen	Recombinant human Parathyroid hormone protein
Conjugation	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300; Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-9677 / sc-9678 / sc-9676 / sc-28922 / sc-28922
Preparation and Storage	Shipped at 4 degree C. Upon delivery, aliquot and store at -20 degree C or -80 degree C.

IMAGES



IHC (Immunohistochemistry)
IHC image of AAA229512 diluted at 1:700 and staining in paraffin-embedded human pancreatic tissue performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



IHC (Immunohistochemistry)
IHC image of AAA229512 diluted at 1:700 and staining in paraffin-embedded human prostate cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.

ADDITIONAL INFORMATION

Related Product Information for anti-PTH antibody	PTH elevates calcium level by dissolving the salts in bone and preventing their renal excretion. Stimulates [1-14C]-2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblastic cells. This protein is a hormone secreted by parathyroid cells. This hormone elevates blood Ca ²⁺ level by dissolving the salts in bone and preventing their renal excretion. Defects in this gene are a cause of familial isolated hypoparathyroidism (FIH).
Product Categories/Family for anti-PTH antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	4506267
NCBI GeneID	5741
NCBI Accession #	NP_000306.1
NCBI GenBank Nucleotide #	NM_000315.3
UniProt Accession #	P01270; Q4VB48; Q9UD38
Molecular Weight	12,861 Da
NCBI Official Full Name	parathyroid hormone isoform 1 preproprotein
NCBI Official Synonym Full Names	parathyroid hormone
NCBI Official Symbol	PTH
NCBI Official Synonym Symbols	PTH1
NCBI Protein Information	parathyroid hormone
UniProt Protein Name	Parathyroid hormone
UniProt Gene Name	PTH
UniProt Synonym Gene Names	PTH
UniProt Entry Name	PTHY_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/parathyroid-hormone-antibody-western-blot-elisa/229512>