

Mouse Toll Like Receptor 4 (TLR4) ELISA Kit | TLR4 elisa kit

Mouse Toll Like Receptor 4 (TLR4) ELISA Kit

CATALOG #
AAA22963

REACTIVITY
Mouse


PRODUCT SYNONYMS

Toll Like Receptor 4 (TLR4); Mouse Toll Like Receptor 4 (TLR4) ELISA Kit; CD284; TOLL; HToll; TLR4 elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of TLR4. No significant cross-reactivity or interference between TLR4 and analogues was observed.
Sequence Length	839
Assay Type	Quantitative Sandwich
Samples	Mouse tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.054ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level TLR4 were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level TLR4 were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for TLR4 elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to TLR4. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to TLR4. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain TLR4, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of TLR4 in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of TLR4 in mouse tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	19924149
NCBI GeneID	7099
NCBI Accession #	NP_612564.1
NCBI GenBank Nucleotide #	NM_138554.4
UniProt Accession #	O00206; Q5VZI8; Q5VZI9; Q9UK78; Q9UM57; A8K1Y8; A9XLP9; A9XLQ0; A9XLQ1; B4E194; D1CS52; D1CS53
Molecular Weight	73,301 Da
NCBI Official Full Name	toll-like receptor 4 isoform A
NCBI Official Synonym Full Names	toll like receptor 4
NCBI Official Symbol	TLR4
NCBI Official Synonym Symbols	TOLL; CD284; TLR-4; ARMD10
NCBI Protein Information	toll-like receptor 4
UniProt Protein Name	Toll-like receptor 4
UniProt Gene Name	TLR4

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.