

Human Involucrin (iNV) ELISA Kit | iNV elisa kit

Human Involucrin (iNV) ELISA Kit

CATALOG #
AAA22999

REACTIVITY
Human


PRODUCT SYNONYMS

Involucrin (iNV); Human Involucrin (iNV) ELISA Kit; iNV; iNV elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of involucrin. No significant cross-reactivity or interference between involucrin and analogues was observed.
Sequence Length	585
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.312-20ng/mL
Sensitivity	< 0.121ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level involucrin were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level involucrin were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for iNV elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to involucrin. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to involucrin. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain involucrin, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of involucrin in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of involucrin in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	44890059
NCBI GeneID	3713
NCBI Accession #	NP_005538.2
NCBI GenBank Nucleotide #	NM_005547.3
UniProt Accession #	P07476; Q5T7P4
Molecular Weight	68,479 Da
NCBI Official Full Name	involucrin
NCBI Official Synonym Full Names	involucrin
NCBI Official Symbol	IVL
NCBI Protein Information	involucrin
UniProt Protein Name	Involucrin
UniProt Gene Name	IVL

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.