

Human Acid Sphingomyelinase (ASM) ELISA Kit | ASM elisa kit

Human Acid Sphingomyelinase (ASM) ELISA Kit

CATALOG #
AAA23001

REACTIVITY
Human


PRODUCT SYNONYMS

Acid Sphingomyelinase (ASM); Human Acid Sphingomyelinase (ASM) ELISA Kit; SMPD1; NPD; SMase; aSMase; Sphingomyelin Phosphodiesterase 1; Acid Lysosomal; Simply Sphingomyelinase; ASM elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Human
Specificity	This assay has high sensitivity and excellent specificity for detection of ASM. No significant cross-reactivity or interference between ASM and analogues was observed.
Sequence Length	628
Assay Type	Quantitative Sandwich
Samples	Human serum, plasma, tissue homogenates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.056ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level ASM were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level ASM were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for ASM elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to ASM. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to ASM. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain ASM, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of ASM in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of ASM in human serum, plasma, tissue homogenates and other biological fluids.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	179095
NCBI GeneID	6609
UniProt Accession #	P17405; P17406; Q13811; Q16837; Q16841; A8K8M3; E9PKS3
Molecular Weight	69,624 Da
NCBI Official Full Name	acid sphingomyelinase
NCBI Official Synonym Full Names	sphingomyelin phosphodiesterase 1
NCBI Official Symbol	SMPD1
NCBI Official Synonym Symbols	ASM; NPD; ASMASE
NCBI Protein Information	sphingomyelin phosphodiesterase
UniProt Protein Name	Sphingomyelin phosphodiesterase
UniProt Gene Name	SMPD1
UniProt Synonym Gene Names	ASM; aSMase

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/human-elisa-kits/acid-sphingomyelinase-asm/23001>