

Rabbit anti-Human MED1 Polyclonal Antibody | anti-MED1 antibody

MED1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA231980	Rabbit	Human	Immunohistochemistry, ELISA


PRODUCT SYNONYMS

MED1; Polyclonal Antibody; MED1 Antibody; Mediator of RNA polymerase II transcription subunit 1; Activator-recruited cofactor 205 kDa component; ARC205; Mediator complex subunit 1; Peroxisome proliferator-activated; CRSP1; CRSP200; DRIP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A; TRAP220; TRIP2; anti-MED1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	1581
Applicable Applications for anti-MED1 antibody	IHC (Immunohistochemistry), ELISA
Species	Human
Immunogen	Recombinant human Mediator of RNA polymerase II transcription subunit 1 protein
Conjugate	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-5335 / sc-5334 / sc-8998
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human thyroid tissue using AAA231980 at dilution of 1:100



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human placenta tissue using AAA231980 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-MED1 antibody	Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (PubMed:10406464, PubMed:11867769, PubMed:12037571, PubMed:12218053, PubMed:12556447, PubMed:14636573, PubMed:15340084, PubMed:15471764, PubMed:15989967, PubMed:16574658, PubMed:9653119). Acts as a coactivator for GATA1-mediated transcriptional activation during erythroid differentiation of K562 erythroleukemia cells (PubMed:24245781).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	28559039
NCBI GeneID	5469
NCBI Accession #	NP_004765.2
NCBI GenBank Nucleotide #	NM_004774.3
UniProt Accession #	Q15648; O43810; O75447; Q6P9H7; Q6PK58; Q9HD39; A2RRQ6
Molecular Weight	61,563 Da
NCBI Official Full Name	mediator of RNA polymerase II transcription subunit 1
NCBI Official Synonym Full Names	mediator complex subunit 1
NCBI Official Symbol	MED1
NCBI Official Synonym Symbols	PBP; CRSP1; RB18A; TRIP2; PPARBP; CRSP200; DRIP205; DRIP230; PPARGBP; TRAP220
NCBI Protein Information	mediator of RNA polymerase II transcription subunit 1
UniProt Protein Name	Mediator of RNA polymerase II transcription subunit 1
UniProt Gene Name	MED1
UniProt Synonym Gene Names	ARC205; CRSP1; CRSP200; DRIP205; DRIP230; PBP; PPARBP; PPARGBP; RB18A; TRAP220; TRIP2; ARC205; PBP; PPAR-binding protein; Trap220; TR-interacting protein 2; TRIP-2
UniProt Entry Name	MED1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

