

Rabbit anti-Human CALCOCO1 Polyclonal Antibody | anti-CALCOCO1 antibody

CALCOCO1 Antibody

CATALOG # AAA232032	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

CALCOCO1; Polyclonal Antibody; CALCOCO1 Antibody; Calcium-binding and coiled-coil domain-containing protein 1; Calphoglin; Coiled-coil coactivator protein; Sarcoma antigen NY-SAR-3; KIAA1536; PP13275; UNQ2436/PRO4996; anti-CALCOCO1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	606
Applicable Applications for anti-CALCOCO1 antibody	IHC (Immunohistochemistry), ELISA
Species	Human
Immunogen	Recombinant human Calcium-binding and coiled-coil domain-containing protein 1 protein (C-230AA)
Conjugate	Non-conjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-161423 / sc-161424
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human liver cancer using AAA232032 at dilution 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human testis tissue using AAA232032 at dilution 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-CALCOCO1 antibody	Functions as a coactivator for aryl hydrocarbon and nuclear receptors (NR). Recruited to promoters through its contact with the N-terminal basic helix-loop-helix-Per-Arnt-Sim (PAS) domain of transcription factors or coactivators, such as NCOA2. During ER-activation acts synergistically in combination with other NCOA2-binding proteins, such as EP300, CREBBP and CARM1. Involved in the transcriptional activation of target genes in the Wnt/CTNNB1 pathway. Functions as a secondary coactivator in LEF1-mediated transcriptional activation via its interaction with CTNNB1. Coactivator function for nuclear receptors and LEF1/CTNNB1 involves differential utilization of two different activation regions (By similarity). In association with CCAR1 enhances GATA1- and MED1-mediated transcriptional activation from the gamma-globin promoter during erythroid differentiation of K562 erythroleukemia cells (PubMed:24245781).
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	219521893
NCBI GeneID	57658
NCBI Accession #	NP_001137154.1
NCBI GenBank Nucleotide #	NM_001143682.1
UniProt Accession #	Q9P1Z2; Q6FI59; Q71RC3; Q86WF8; Q96JU3; Q9H090; B3KVA8
Molecular Weight	67,542 Da
NCBI Official Full Name	calcium-binding and coiled-coil domain-containing protein 1 isoform 2
NCBI Official Synonym Full Names	calcium binding and coiled-coil domain 1
NCBI Official Symbol	CALCOCO1
NCBI Official Synonym Symbols	Cocoa; PP13275; calphoglin
NCBI Protein Information	calcium-binding and coiled-coil domain-containing protein 1
UniProt Protein Name	Calcium-binding and coiled-coil domain-containing protein 1
UniProt Gene Name	CALCOCO1
UniProt Synonym Gene Names	KIAA1536
UniProt Entry Name	CACO1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/calcoco1-antibody-immunohistochemistry-elisa/232032>