

Rabbit anti-Human EPHB4 Polyclonal Antibody | anti-EPHB4 antibody

EPHB4 Antibody

CATALOG # AAA232103	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

EPHB4; Polyclonal Antibody; EPHB4 Antibody; Ephrin type-B receptor 4; Hepatoma transmembrane kinase; Tyrosine-protein kinase TYRO11; HTK; MYK1; TYRO11; anti-EPHB4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid
Sequence Length	987
Applicable Applications for anti-EPHB4 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Species	Human
Immunogen	Recombinant human Ephrin type-B receptor 4 protein (300-540AA)
Conjugate	Non-conjugated
Buffer	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-7285 / sc-7284-R / sc-130753 / sc-5536
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.


IMAGES



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human kidney tissue AAA232103 at dilution 1:100



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human liver tissue AAA232103 at dilution 1:100



WB (Western Blot)
 Western blotAll lanes: EPHB4 antibody at 2.71ug/mlLane 1: A549 whole cell lysateLane 2: 293T whole cell lysateLane 3: Hela whole cell lysateSecondaryGoat polyclonal to rabbit IgG at 1/10000 dilutionPredicted band size: 109,56,34,46 kDaObserved band size: 108 kDa

ADDITIONAL INFORMATION

Related Product Information for anti-EPHB4 antibody	Receptor tyrosine kinase which binds promiscuously transmembrane ephrin-B family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Together with its cognate ligand/functional ligand EFNB2 plays a central role in heart morphogenesis and angiogenesis through regulation of cell adhesion and cell migration. EPHB4-mediated forward signaling controls cellular repulsion and segregation form EFNB2-expressing cells. Plays also a role in postnatal blood vessel remodeling, morphogenesis and permeability and is thus important in the context of tumor angiogenesis.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	32528301
NCBI GeneID	2050
NCBI Accession #	NP_004435.3
NCBI GenBank Nucleotide #	NM_004444.4
UniProt Accession #	P54760; Q7Z635; Q9BTA5; Q9BXP0; B5A970; B5A971; B5A972
Molecular Weight	45,195 Da
NCBI Official Full Name	ephrin type-B receptor 4
NCBI Official Synonym Full Names	EPH receptor B4
NCBI Official Symbol	EPHB4
NCBI Official Synonym Symbols	HTK; MYK1; TYRO11
NCBI Protein Information	ephrin type-B receptor 4
UniProt Protein Name	Ephrin type-B receptor 4
UniProt Gene Name	EPHB4
UniProt Synonym Gene Names	HTK; MYK1; TYRO11
UniProt Entry Name	EPHB4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

