

Rabbit anti-Human SMARCA4 Polyclonal Antibody | anti-SMARCA4 antibody

SMARCA4 Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232306	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA

PRODUCT SYNONYMS


SMARCA4; Polyclonal Antibody; SMARCA4 Antibody; Transcription activator BRG1; ATP-dependent helicase SMARCA4; BRG1-associated factor 190A; BAF190A; Mitotic growth and transcription activator; Prote; BRG1; SNF2B; SNF2L4; anti-SMARCA4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	1647
Applicable Applications for anti-SMARCA4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Species	Human
Immunogen	Recombinant human Transcription activator BRG1 protein (808-1092AA)
Conjugate	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-8749 / sc-12520 / sc-10768
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.



IF (Immunofluorescence)
Immunofluorescent analysis of HepG-2 cells using AAA232306 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)
Immunohistochemistry of paraffin-embedded human lung cancer using AAA232306 at dilution 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-SMARCA4 antibody	Transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. Component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. In resting neurons, transcription of the c-FOS promoter is inhibited by BRG1-dependent recruitment of a phospho-RB1-HDAC repressor complex. Upon calcium influx, RB1 is dephosphorylated by calcineurin, which leads to release of the repressor complex. At the same time, there is increased recruitment of CREBBP to the promoter by a CREST-dependent mechanism, which leads to transcriptional activation. The CREST-BRG1 complex also binds to the NR2B promoter, and activity-dependent induction of NR2B expression involves a release of HDAC1 and recruitment of CREBBP. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth. SMARCA4/BAF190A may promote neural stem cell self-renewal/proliferation by enhancing Notch-dependent proliferative signals, while concurrently making the neural stem cell insensitive to SHH-dependent differentiating cues (By similarity). Acts as a corepressor of ZEB1 to regulate E-cadherin transcription and is required for induction of epithelial-mesenchymal transition (EMT) by ZEB1.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	192807312
NCBI GeneID	6597
NCBI Accession #	NP_001122316.1
NCBI GenBank Nucleotide #	NM_001128844.1
UniProt Accession #	P51532; O95052; Q9HBD3; B1A8Z4; B1A8Z5; B1A8Z6; B1A8Z7; E9PBR8
Molecular Weight	181,261 Da
NCBI Official Full Name	transcription activator BRG1 isoform B
NCBI Official Synonym Full Names	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4
NCBI Official Symbol	SMARCA4
NCBI Official Synonym Symbols	BRG1; CSS4; SNF2; SWI2; MRD16; RTPS2; BAF190; SNF2L4; SNF2LB; hSNF2b; BAF190A

NCBI Protein Information	transcription activator BRG1
UniProt Protein Name	Transcription activator BRG1
UniProt Gene Name	SMARCA4
UniProt Synonym Gene Names	BAF190A; BRG1; SNF2B; SNF2L4; BAF190A
UniProt Entry Name	SMCA4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/smarca4-antibody-immunofluorescence-immunohistochemistry-elisa/232306>