

Rabbit anti-Human, Mouse BTK Polyclonal Antibody | anti-BTK antibody

BTK Antibody

CATALOG # AAA232411	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunofluorescence, Western Blot, ELISA
-------------------------------	-----------------------	-----------------------------------	--


PRODUCT SYNONYMS

BTK; Polyclonal Antibody; BTK Antibody; Tyrosine-protein kinase BTK; Agammaglobulinemia tyrosine kinase; ATK; B-cell progenitor kinase; BPK; Bruton tyrosine kinase; AGMX1; anti-BTK antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	659
Applicable Applications for anti-BTK antibody	IF (Immunofluorescence), WB (Western Blot), ELISA
Immunogen	Recombinant Human Tyrosine-protein kinase BTK protein (254-443AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-1107 / sc-1108 / sc-1696
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescent analysis of HeLa cells using AAA232411 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human spleen tissue using AAA232411 at dilution of 1:100



WB (Western Blot)

Western Blot Positive WB detected in: K562 whole cell lysate, HL60 whole cell lysate, Mouse heart tissue, Mouse liver tissue, Mouse brain tissue. All lanes: BTK antibody at 3.5 µg/ml. Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 77,80 kDa. Observed band size: 77 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-BTK antibody

Non-receptor tyrosine kinase indispensable for B lymphocyte development, differentiation and signaling. Binding of antigen to the B-cell antigen receptor (BCR) triggers signaling that ultimately leads to B-cell activation. After BCR engagement and activation at the plasma membrane, phosphorylates PLCG2 at several sites, igniting the downstream signaling pathway through calcium mobilization, followed by activation of the protein kinase C (PKC) family members. PLCG2 phosphorylation is performed in close cooperation with the adapter protein B-cell linker protein BLNK. BTK acts as a platform to bring together a diverse array of signaling proteins and is implicated in cytokine receptor signaling pathways. Plays an important role in the function of immune cells of innate as well as adaptive immunity, as a component of the Toll-like receptors (TLR) pathway. The TLR pathway acts as a primary surveillance system for the detection of pathogens and are crucial to the activation of host defense. Especially, is a critical molecule in regulating TLR9 activation in splenic B-cells. Within the TLR pathway, induces tyrosine phosphorylation of TIRAP which leads to TIRAP degradation. BTK plays also a critical role in transcription regulation. Induces the activity of NF-kappa-B, which is involved in regulating the expression of hundreds of genes. BTK is involved on the signaling pathway linking TLR8 and TLR9 to NF-kappa-B. Transiently phosphorylates transcription factor GTF2I on tyrosine residues in response to BCR. GTF2I then translocates to the nucleus to bind regulatory enhancer elements to modulate gene expression. ARID3A and NFAT are other transcriptional target of BTK. BTK is required for the formation of functional ARID3A DNA-binding complexes. There is however no evidence that BTK itself binds directly to DNA. BTK has a dual role in the regulation of apoptosis.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557377
NCBI GeneID	695
NCBI Accession #	NP_000052.1
NCBI GenBank Nucleotide #	NM_000061.2
UniProt Accession #	Q06187 ; Q32ML5 ; B2RAW1
Molecular Weight	79,937 Da
NCBI Official Full Name	tyrosine-protein kinase BTK isoform 1
NCBI Official Synonym Full Names	Bruton tyrosine kinase
NCBI Official Symbol	BTK
NCBI Official Synonym Symbols	AT; ATK; BPK; XLA; IMD1; AGMX1; PSCTK1
NCBI Protein Information	tyrosine-protein kinase BTK
UniProt Protein Name	Tyrosine-protein kinase BTK

UniProt Gene Name	BTK
UniProt Synonym Gene Names	AGMX1; ATK; BPK; ATK; BPK
UniProt Entry Name	BTK_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/btk-antibody-immunofluorescence-western-blot-elisa/232411>