

Rabbit anti-Human EIF2S1 Polyclonal Antibody | anti-EIF2S1 antibody

EIF2S1 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232467	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

EIF2S1; Polyclonal Antibody; EIF2S1 Antibody; Eukaryotic translation initiation factor 2 subunit 1; Eukaryotic translation initiation factor 2 subunit alpha; eIF-2-alpha; eIF-2A; eIF-2alpha; EIF2A; anti-EIF2S1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	315
Applicable Applications for anti-EIF2S1 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Eukaryotic translation initiation factor 2 subunit 1 protein (276-315AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-11386 / sc-30882
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescent analysis of MCF-7 cells using AAA232467 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human brain tissue using AAA232467 at dilution of 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon cancer using AAA232467 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-EIF2S1 antibody

Functions in the early steps of protein synthesis by forming a ternary complex with GTP and initiator tRNA. This complex binds to a 40S ribosomal subunit, followed by mRNA binding to form a 43S preinitiation complex. Junction of the 60S ribosomal subunit to form the 80S initiation complex is preceded by hydrolysis of the GTP bound to eIF-2 and release of an eIF-2-GDP binary complex. In order for eIF-2 to recycle and catalyze another round of initiation, the GDP bound to eIF-2 must exchange with GTP by way of a reaction catalyzed by eIF-2B.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758256
NCBI GeneID	1965
NCBI Accession #	NP_004085.1
NCBI GenBank Nucleotide #	NM_004094.4
UniProt Accession #	P05198
Molecular Weight	36,112 Da
NCBI Official Full Name	eukaryotic translation initiation factor 2 subunit 1
UniProt Protein Name	Eukaryotic translation initiation factor 2 subunit 1
UniProt Gene Name	EIF2S1
UniProt Synonym Gene Names	EIF2A; eIF-2-alpha; eIF-2A; eIF-2alpha
UniProt Entry Name	IF2A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://www.aaabiotech.com/polyclonal/eif2s1-antibody-immunofluorescence-immunohistochemistry-elisa/232467>