

Rabbit anti-Human ERBB4 Polyclonal Antibody | anti-ERBB4 antibody

ERBB4 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232479	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

ERBB4; Polyclonal Antibody; ERBB4 Antibody; Receptor tyrosine-protein kinase erbB-4; Proto-oncogene-like protein c-ErbB-4; Tyrosine kinase-type cell surface receptor HER4; p180erbB4; HER4; anti-ERBB4 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	1292
Applicable Applications for anti-ERBB4 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Receptor tyrosine-protein kinase erbB-4 protein (1027-1204AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-31151 / sc-283
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES


IF (Immunofluorescence)

Immunofluorescent analysis of HepG2 cells using AAA232479 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)


IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human placenta tissue using AAA232479 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-ERBB4 antibody	Tyrosine-protein kinase that plays an essential role as cell surface receptor for neuregulins and EGF family members and regulates development of the heart, the central nervous system and the mammary gland, gene transcription, cell proliferation, differentiation, migration and apoptosis. Required for normal cardiac muscle differentiation during embryonic development, and for postnatal cardiomyocyte proliferation. Required for normal development of the embryonic central nervous system, especially for normal neural crest cell migration and normal axon guidance. Required for mammary gland differentiation, induction of milk proteins and lactation. Acts as cell-surface receptor for the neuregulins NRG1, NRG2, NRG3 and NRG4 and the EGF family members BTC, EREG and HBEGF. Ligand binding triggers receptor dimerization and autophosphorylation at specific tyrosine residues that then serve as binding sites for scaffold proteins and effectors. Ligand specificity and signaling is modulated by alternative splicing, proteolytic processing, and by the formation of heterodimers with other ERBB family members, thereby creating multiple combinations of intracellular phosphotyrosines that trigger ligand- and context-specific cellular responses. Mediates phosphorylation of SHC1 and activation of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1. Isoform JM-A CYT-1 and isoform JM-B CYT-1 phosphorylate PIK3R1, leading to the activation of phosphatidylinositol 3-kinase and AKT1 and protect cells against apoptosis. Isoform JM-A CYT-1 and isoform JM-B CYT-1 mediate reorganization of the actin cytoskeleton and promote cell migration in response to NRG1. Isoform JM-A CYT-2 and isoform JM-B CYT-2 lack the phosphotyrosine that mediates interaction with PIK3R1, and hence do not phosphorylate PIK3R1, do not protect cells against apoptosis, and do not promote reorganization of the actin cytoskeleton and cell migration. Proteolytic processing of isoform JM-A CYT-1 and isoform JM-A CYT-2 gives rise to the corresponding soluble intracellular domains (4ICD) that translocate to the nucleus, promote nuclear import of STAT5A, activation of STAT5A, mammary epithelium differentiation, cell proliferation and activation of gene expression. The ERBB4 soluble intracellular domains (4ICD) colocalize with STAT5A at the CSN2 promoter to regulate transcription of milk proteins during lactation. The ERBB4 soluble intracellular domains can also translocate to mitochondria and promote apoptosis.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	110825958
NCBI GeneID	2066
NCBI Accession #	NP_001036064.1
NCBI GenBank Nucleotide #	NM_001042599.1
UniProt Accession #	Q15303; Q2M1W1; Q59EW4; B7ZLD7; B7ZLE2; B7ZLE3
Molecular Weight	143,968 Da
NCBI Official Full Name	receptor tyrosine-protein kinase erbB-4 isoform JM-a/CVT-2
NCBI Official Synonym Full Names	erb-b2 receptor tyrosine kinase 4
NCBI Official Symbol	ERBB4
NCBI Official Synonym Symbols	HER4; ALS19; p180erbB4
NCBI Protein Information	receptor tyrosine-protein kinase erbB-4
UniProt Protein Name	Receptor tyrosine-protein kinase erbB-4
UniProt Gene Name	ERBB4
UniProt Synonym Gene Names	HER4; 4ICD; E4ICD
UniProt Entry Name	ERBB4_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.