

Rabbit anti-Human MAPK10 Polyclonal Antibody | anti-MAPK10 antibody

MAPK10 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232576	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

MAPK10; Polyclonal Antibody; MAPK10 Antibody; Mitogen-activated protein kinase 10; MAP kinase p49 3F12; Stress-activated protein kinase 1b; SAPK1b; Stress-activated protein kinase JNK3; c-Jun N-terminal; JNK3; JNK3A; PRKM10; SAPK1B; anti-MAPK10 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	464
Applicable Applications for anti-MAPK10 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant human Mitogen-activated protein kinase 10 protein (133-316AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-46014 / sc-46015
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescent analysis of HepG2 cells using AAA232576 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human testis tissue using AAA232576 at dilution of 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human brain tissue using AAA232576 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-MAPK10 antibody

Serine/threonine-protein kinase involved in various processes such as neuronal proliferation, differentiation, migration and programmed cell death. Extracellular stimuli such as proinflammatory cytokines or physical stress stimulate the stress-activated protein kinase/c-Jun N-terminal kinase (SAP/JNK) signaling pathway. In this cascade, two dual specificity kinases MAP2K4/MKK4 and MAP2K7/MKK7 phosphorylate and activate MAPK10/JNK3. In turn, MAPK10/JNK3 phosphorylates a number of transcription factors, primarily components of AP-1 such as JUN and ATF2 and thus regulates AP-1 transcriptional activity. Plays regulatory roles in the signaling pathways during neuronal apoptosis. Phosphorylates the neuronal microtubule regulator STMN2. Acts in the regulation of the beta-amyloid precursor protein/APP signaling during neuronal differentiation by phosphorylating APP. Participates also in neurite growth in spiral ganglion neurons. Phosphorylates the CLOCK-ARNTL/BMAL1 heterodimer and plays a role in the photic regulation of the circadian clock (PubMed: 22441692).

NCBI AND UNIPROT INFORMATION

NCBI GI #	969536252
NCBI GeneID	5602
NCBI Accession #	NP_001304996.1
NCBI GenBank Nucleotide #	NM_001318067.1
UniProt Accession #	P53779; Q15707; Q49AP1; A6NFS3; A6NG28; B3KQ94
Molecular Weight	31,933 Da
NCBI Official Full Name	mitogen-activated protein kinase 10 isoform 5
NCBI Official Synonym Full Names	mitogen-activated protein kinase 10
NCBI Official Symbol	MAPK10
NCBI Official Synonym Symbols	JNK3; JNK3A; PRKM10; SAPK1b; p493F12; p54bSAPK
NCBI Protein Information	mitogen-activated protein kinase 10
UniProt Protein Name	Mitogen-activated protein kinase 10
UniProt Gene Name	MAPK10
UniProt Synonym Gene Names	JNK3; JNK3A; PRKM10; SAPK1B; MAP kinase 10; MAPK 10; SAPK1b
UniProt Entry Name	MK10_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.