

Rabbit anti-Human TNFSF11 Polyclonal Antibody | anti-TNFSF11 antibody

TNFSF11 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232714	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

TNFSF11; Polyclonal Antibody; TNFSF11 Antibody; Tumor necrosis factor ligand superfamily member 11; Osteoclast differentiation factor; ODF; Osteoprotegerin ligand; OPGL; Receptor activator of nuclear factor kappa-B ligand; RANKL; TRANCE; anti-TNFSF11 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	317
Applicable Applications for anti-TNFSF11 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Tumor necrosis factor ligand superfamily member 11 protein (73-306AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-7627 / sc-7628 / sc-9073
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescent analysis of HepG2 cells using AAA232714 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon cancer using AAA232714 at dilution of 1:100



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human gastric cancer using AAA232714 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-TNFSF11 antibody

Cytokine that binds to TNFRSF11B/OPG and to TNFRSF11A/RANK. Osteoclast differentiation and activation factor. Augments the ability of dendritic cells to stimulate naive T-cell proliferation. May be an important regulator of interactions between T-cells and dendritic cells and may play a role in the regulation of the T-cell-dependent immune response. May also play an important role in enhanced bone-resorption in humoral hypercalcemia of malignancy (PubMed: 22664871). Induces osteoclastogenesis by activating multiple signaling pathways in osteoclast precursor cells, chief among which is induction of long lasting oscillations in the intracellular concentration of Ca (2+) resulting in the activation of NFATC1, which translocates to the nucleus and induces osteoclast-specific gene transcription to allow differentiation of osteoclasts. During osteoclast differentiation, in a TMEM64 and ATP2A2-dependent manner induces activation of CREB1 and mitochondrial ROS generation necessary for proper osteoclast generation (By similarity).

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507595
NCBI GeneID	8600
NCBI Accession #	NP_003692.1
NCBI GenBank Nucleotide #	NM_003701.3
UniProt Accession #	O14788; O14723; Q96Q17; Q9P2Q3
Molecular Weight	30,523 Da
NCBI Official Full Name	tumor necrosis factor ligand superfamily member 11 isoform 1
NCBI Official Synonym Full Names	tumor necrosis factor superfamily member 11
NCBI Official Symbol	TNFSF11
NCBI Official Synonym Symbols	ODF; OPGL; sOdf; CD254; OPTB2; RANKL; TNLG6B; TRANCE; hRANKL2
NCBI Protein Information	tumor necrosis factor ligand superfamily member 11
UniProt Protein Name	Tumor necrosis factor ligand superfamily member 11
UniProt Gene Name	TNFSF11
UniProt Synonym Gene Names	OPGL; RANKL; TRANCE; ODF; OPGL; RANKL; TRANCE
UniProt Entry Name	TNF11_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.