

Rabbit anti-Human FLT1 Polyclonal Antibody | anti-FLT1 antibody

FLT1 Antibody, HRP conjugated

CATALOG # AAA232745	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	-----------------------	----------------------------	------------------------------

PRODUCT SYNONYMS

FLT1; Polyclonal Antibody; FLT1 Antibody; HRP conjugated; Vascular endothelial growth factor receptor 1; Fms-like tyrosine kinase 1; FLT-1; Tyrosine-protein kinase FRT; Tyrosine-protein kinase receptor FLT; FLT; Vascular perm; FRT; VEGFR1; anti-FLT1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Applicable Applications for anti-FLT1 antibody	ELISA
Conjugation	HRP
Research Area	Cardiovascular
Immunogen	Recombinant human Vascular endothelial growth factor receptor 1 protein (1048-1328AA)
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-31173 / sc-9029 / sc-316
Preparation and Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

ADDITIONAL INFORMATION

Related Product Information for anti-FLT1 antibody	Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFA, VEGFB and PGF, and plays an essential role in the development of embryonic vasculature, the regulation of angiogenesis, cell survival, cell migration, macrophage function, chemotaxis, and cancer cell invasion. May play an essential role as a negative regulator of embryonic angiogenesis by inhibiting excessive proliferation of endothelial cells. Can promote endothelial cell proliferation, survival and angiogenesis in adulthood. Its function in promoting cell proliferation seems to be cell-type specific. Promotes PGF-mediated proliferation of endothelial cells, proliferation of some types of cancer cells, but does not promote proliferation of normal fibroblasts (in vitro). Has very high affinity for VEGFA and relatively low protein kinase activity; may function as a negative regulator of VEGFA signaling by limiting the amount of free VEGFA and preventing its binding to KDR. Likewise, isoforms lacking a transmembrane domain, such as isoform 2, isoform 3 and isoform 4, may function as decoy receptors for VEGFA. Modulates KDR signaling by forming heterodimers with KDR. Ligand binding leads to the activation of several signaling cascades. Activation of PLCG leads to the production of the cellular signaling molecules diacylglycerol and inositol 1, 4, 5-trisphosphate and the activation of protein kinase C. Mediates phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, leading to activation of phosphatidylinositol kinase and the downstream signaling pathway. Mediates activation of MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling pathway, as well as of the AKT1 signaling pathway. Phosphorylates SRC and YES1, and may also phosphorylate CBL. Isoform 1 phosphorylates PLCG. Promotes phosphorylation of AKT1 at 'Ser-473'. Promotes phosphorylation of PTK2/FAK1. Isoform 7 has a truncated kinase domain; it increases phosphorylation of SRC at 'Tyr-418' by unknown means and promotes tumor cell invasion.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	229892220
NCBI GeneID	2321
NCBI Accession #	NP_001153392.1
NCBI GenBank Nucleotide #	NM_001159920.1
UniProt Accession #	P17948; O60722; A3E342; A3E344; A8KA71; B0LPF1; B2BF46; B2BF47; B2BF48; B3FR89; B5A923; F5H5L6
NCBI Official Full Name	vascular endothelial growth factor receptor 1 isoform 2
NCBI Official Synonym Full Names	fms related tyrosine kinase 1
NCBI Official Symbol	FLT1
NCBI Official Synonym Symbols	FLT; FLT-1; VEGFR1; VEGFR-1
NCBI Protein Information	vascular endothelial growth factor receptor 1
UniProt Protein Name	Vascular endothelial growth factor receptor 1
UniProt Gene Name	FLT1
UniProt Synonym Gene Names	FLT; FRT; VEGFR1; VEGFR-1; FLT-1; FLT
UniProt Entry Name	VGFR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.