

Rabbit anti-Human FLT4 Polyclonal Antibody | anti-FLT4 antibody

FLT4 Antibody, HRP conjugated

CATALOG # AAA232749	HOST Rabbit	REACTIVITY Human	APPLICATIONS ELISA
-------------------------------	-----------------------	----------------------------	------------------------------

PRODUCT SYNONYMS

FLT4; Polyclonal Antibody; FLT4 Antibody; HRP conjugated; Vascular endothelial growth factor receptor 3; VEGFR-3; Fms-like tyrosine kinase 4; FLT-4; Tyrosine-protein kinase receptor FLT4; VEGFR3; anti-FLT4 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Applicable Applications for anti-FLT4 antibody	ELISA
Conjugation	HRP
Research Area	Cardiovascular
Immunogen	Recombinant Human Vascular endothelial growth factor receptor 3 protein (1112-1329AA)
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-20734 / sc-321 / sc-637
Preparation and Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

ADDITIONAL INFORMATION

Related Product Information for anti-FLT4 antibody	Tyrosine-protein kinase that acts as a cell-surface receptor for VEGFC and VEGFD, and plays an essential role in adult lymphangiogenesis and in the development of the vascular network and the cardiovascular system during embryonic development. Promotes proliferation, survival and migration of endothelial cells, and regulates angiogenic sprouting. Signaling by activated FLT4 leads to enhanced production of VEGFC, and to a lesser degree VEGFA, thereby creating a positive feedback loop that enhances FLT4 signaling. Modulates KDR signaling by forming heterodimers. The secreted isoform 3 may function as a decoy receptor for VEGFC and/or VEGFD and play an important role as a negative regulator of VEGFC-mediated lymphangiogenesis and angiogenesis. Binding of vascular growth factors to isoform 1 or isoform 2 leads to the activation of several signaling cascades; isoform 2 seems to be less efficient in signal transduction, because it has a truncated C-terminus and therefore lacks several phosphorylation sites. Mediates activation of the MAPK1/ERK2, MAPK3/ERK1 signaling pathway, of MAPK8 and the JUN signaling pathway, and of the AKT1 signaling pathway. Phosphorylates SHC1. Mediates phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase. Promotes phosphorylation of MAPK8 at 'Thr-183' and 'Tyr-185', and of AKT1 at 'Ser-473'.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	103472027
NCBI GeneID	2324
NCBI Accession #	NP_002011.2
NCBI GenBank Nucleotide #	NM_002020.4
UniProt Accession #	P35916; Q16067; Q86W07; Q86W08; A8K6L4; B5A926; AAB23636
NCBI Official Full Name	vascular endothelial growth factor receptor 3 isoform 2
NCBI Official Synonym Full Names	fms related tyrosine kinase 4
NCBI Official Symbol	FLT4
NCBI Official Synonym Symbols	PCL; FLT-4; FLT41; LMPH1A; VEGFR3; VEGFR-3
NCBI Protein Information	vascular endothelial growth factor receptor 3
UniProt Protein Name	Vascular endothelial growth factor receptor 3
UniProt Gene Name	FLT4
UniProt Synonym Gene Names	VEGFR3; VEGFR-3; FLT-4
UniProt Entry Name	VGFR3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/flt4-antibody-elisa/232749>