

Rabbit anti-Human, Mouse NGFR Polyclonal Antibody | anti-NGFR antibody

NGFR Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232757	Rabbit	Human, Mouse	Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

NGFR; Polyclonal Antibody; NGFR Antibody; Tumor necrosis factor receptor superfamily member 16; Gp80-LNGFR; Low affinity neurotrophin receptor p75NTR; Low-affinity nerve growth factor receptor; NGF receptor; p75 ICD; CD271; TNFRSF16; anti-NGFR antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	427
Applicable Applications for anti-NGFR antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant Human Tumor necrosis factor receptor superfamily member 16 protein (277-391AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-5634 / sc-6188 / sc-6189 / sc-8317
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IHC (Immunohistochemistry)
IHC image of AAA232757 diluted at 1:400 and staining in paraffin-embedded human glioma cancer performed on a Leica Bond™ system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4 degree C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



WB (Western Blot)
Western BlotPositive WB detected in: Mouse liver tissueAll lanes: NGFR antibody at 2.7ug/mlSecondaryGoat polyclonal to rabbit IgG at 1/50000 dilutionPredicted band size: 46,36 KDaObserved band size: 46 KDa

ADDITIONAL INFORMATION

Related Product Information for anti-NGFR antibody	Plays a role in the regulation of the translocation of GLUT4 to the cell surface in adipocytes and skeletal muscle cells in response to insulin, probably by regulating RAB31 activity, and thereby contributes to the regulation of insulin-dependent glucose uptake (By similarity). Low affinity receptor which can bind to NGF, BDNF, NT-3, and NT-4. Can mediate cell survival as well as cell death of neural cells. Necessary for the circadian oscillation of the clock genes ARNTL/BMAL1, PER1, PER2 and NR1D1 in the suprachiasmatic nucleus (SCN) of the brain and in liver and of the genes involved in glucose and lipid metabolism in the liver.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4505393
NCBI GeneID	4804
NCBI Accession #	NP_002498.1
NCBI GenBank Nucleotide #	NM_002507.3
UniProt Accession #	P08138; B2R961; B4E096
Molecular Weight	35,767 Da
NCBI Official Full Name	tumor necrosis factor receptor superfamily member 16
NCBI Official Synonym Full Names	nerve growth factor receptor
NCBI Official Symbol	NGFR
NCBI Official Synonym Symbols	CD271; p75NTR; TNFRSF16; p75(NTR); Gp80-LNGFR
NCBI Protein Information	tumor necrosis factor receptor superfamily member 16
UniProt Protein Name	Tumor necrosis factor receptor superfamily member 16
UniProt Gene Name	NGFR
UniProt Synonym Gene Names	TNFRSF16; NGF receptor
UniProt Entry Name	TNR16_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

