

Rabbit anti-Human, Mouse MAPT Polyclonal Antibody | anti-MAPT antibody

MAPT Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA232760	Rabbit	Human, Mouse	Immunofluorescence, Immunohistochemistry, Western Blot, ELISA


PRODUCT SYNONYMS

MAPT; Polyclonal Antibody; MAPT Antibody; Microtubule-associated protein tau; Neurofibrillary tangle protein; Paired helical filament-tau; PHF-tau; MAPTL; MTBT1; TAU; anti-MAPT antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	776
Applicable Applications for anti-MAPT antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant human Microtubule-associated protein tau protein (33-166AA)
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, PH 7.4
Conjugation	Non-conjugated
Santa Cruz Alternative	Potential replacement for Santa Cruz Biotechnology antibody catalog# sc-1995 / sc-1996 / sc-5587
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES




IHC (Immunohistochemistry)

Immunofluorescent analysis of HepG2 cells using AAA232760 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human brain tissue using AAA232760 at dilution of 1:100



WB (Western Blot)

Western Blot Positive WB detected in: SH-SY5Y whole cell lysate, Mouse heart tissue. All lanes: MAPT antibody at 3.25 µg/ml. Secondary: Goat polyclonal to rabbit IgG at 1/50000 dilution. Predicted band size: 79, 37, 33, 40, 43, 41, 46, 81 kDa. Observed band size: 79, 50, 36 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-MAPT antibody	Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by TAU/MAPT localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	294862258
NCBI GeneID	4137
NCBI Accession #	NP_001116538.2
NCBI GenBank Nucleotide #	NM_001123066.3
UniProt Accession #	P10636; P18518; Q14799; Q15549; Q15550; Q15551; Q1RMF6; Q53YB1; Q5CZ17; Q5XWF0; Q6QT54; Q9UDJ3
Molecular Weight	80,941 Da
NCBI Official Full Name	microtubule-associated protein tau isoform 6
NCBI Official Synonym Full Names	microtubule associated protein tau
NCBI Official Symbol	MAPT
NCBI Official Synonym Symbols	TAU; MSTD; PPND; DDPAC; MAPTL; MTBT1; MTBT2; FTDP-17; PPP1R103
NCBI Protein Information	microtubule-associated protein tau
UniProt Protein Name	Microtubule-associated protein tau
UniProt Gene Name	MAPT
UniProt Synonym Gene Names	MAPTL; MTBT1; TAU; PHF-tau
UniProt Entry Name	TAU_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.