

Rabbit anti-Human TRAF3 Polyclonal Antibody | anti-TRAF3 antibody

TRAF3 Antibody

CATALOG # AAA233305	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunofluorescence, ELISA
-------------------------------	-----------------------	----------------------------	--


PRODUCT SYNONYMS

TRAF3; Polyclonal Antibody; TRAF3 Antibody; TNF receptor-associated factor 3; CAP-1; CD40 receptor-associated factor 1; CRAFT1; CD40-binding protein; CD40BP; LMP1-associated protein 1; LAP1; CAP1; anti-TRAF3 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	485
Applicable Applications for anti-TRAF3 antibody	IF (Immunofluorescence), ELISA
Immunogen	Recombinant Human TNF receptor-associated factor 3 protein (1-40AA)
Conjugation	Non-conjugated
Preservative	0.03% Proclin 300
Constituents	50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Shipped at 4 degree C. Upon delivery aliquot and store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



WB (Western Blot)

Western Blot Positive WB detected in: HEK293 whole cell lysate All lanes: TRAF3 antibody at 3.4ug/ml Secondary Goat polyclonal to rabbit IgG at 1/50000 dilution Predicted band size: 65,56 KDa Observed band size: 65 KDa



IF (Immunofluorescence)

Immunofluorescent analysis of HeLa cells using AAA233305 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)

ADDITIONAL INFORMATION

Related Product Information for anti-TRAF3 antibody	Regulates pathways leading to the activation of NF-kappa-B and MAP kinases, and plays a central role in the regulation of B-cell survival. Part of signaling pathways leading to the production of cytokines and interferon. Required for normal antibody isotype switching from IgM to IgG. Plays a role T-cell dependent immune responses. Plays a role in the regulation of antiviral responses. Is an essential constituent of several E3 ubiquitin-protein ligase complexes. May have E3 ubiquitin-protein ligase activity and promote 'Lys-63'-linked ubiquitination of target proteins. Inhibits activation of NF-kappa-B in response to LTBR stimulation. Inhibits TRAF2-mediated activation of NF-kappa-B. Down-regulates proteolytic processing of NFKB2, and thereby inhibits non-canonical activation of NF-kappa-B. Promotes ubiquitination and proteasomal degradation of MAP3K14.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	313661483
NCBI GeneID	7187
NCBI Accession #	NP_001186356.1
NCBI GenBank Nucleotide #	NM_001199427.1
UniProt Accession #	Q13114; Q12990; Q13076; Q13947; Q6AZX1; Q9UNL1; B7Z8C4
Molecular Weight	55,306 Da
NCBI Official Full Name	TNF receptor-associated factor 3 isoform 3
NCBI Official Synonym Full Names	TNF receptor associated factor 3
NCBI Official Symbol	TRAF3
NCBI Official Synonym Symbols	CAP1; LAP1; CAP-1; CRAF1; IIAE5; CD40bp
NCBI Protein Information	TNF receptor-associated factor 3
UniProt Protein Name	TNF receptor-associated factor 3
UniProt Gene Name	TRAF3
UniProt Synonym Gene Names	CAP1; CRAF1; CRAF1; CD40BP; LAP1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/traf3-antibody-immunofluorescence-elisa/233305>