

Rabbit anti-Human PKN2 Polyclonal Antibody | anti-PKN2 antibody

PKN2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA233727	Rabbit	Human	Immunofluorescence, Immunohistochemistry, ELISA


PRODUCT SYNONYMS

PKN2; Polyclonal Antibody; PKN2 Antibody; Serine/threonine-protein kinase N2; PKN gamma; Protein kinase C-like 2; Protein-kinase C-related kinase 2; PRK2; PRKCL2; anti-PKN2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Purity/Purification	>95%, Protein G purified
Form/Format	Liquid
Sequence Length	936
Applicable Applications for anti-PKN2 antibody	IF (Immunofluorescence), IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant Human Serine/threonine-protein kinase N2 protein (925-969AA)
Conjugation	Non-conjugated
Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Preparation and Storage	Upon receipt, store at -20 degree C or -80 degree C. Avoid repeated freeze.

IMAGES



IF (Immunofluorescence)

Immunofluorescent analysis of HepG2 Cells using AAA233727 at a dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



IHC (Immunohistochemistry)

Immunohistochemistry of paraffin-embedded human colon cancer using AAA233727 at dilution of 1:100

ADDITIONAL INFORMATION

Related Product Information for anti-PKN2 antibody	PKC-related serine/threonine-protein kinase and Rho/Rac effector protein that participates in specific signal transduction responses in the cell. Plays a role in the regulation of cell cycle progression, actin cytoskeleton assembly, cell migration, cell adhesion, tumor cell invasion and transcription activation signaling processes. Phosphorylates CTTN in hyaluronan-induced astrocytes and hence decreases CTTN ability to associate with filamentous actin. Phosphorylates HDAC5, therefore lead to impair HDAC5 import. Direct RhoA target required for the regulation of the maturation of primordial junctions into apical junction formation in bronchial epithelial cells. Required for G2/M phases of the cell cycle progression and abscission during cytokinesis in a ECT2-dependent manner. Stimulates FYN kinase activity that is required for establishment of skin cell-cell adhesion during keratinocytes differentiation. Regulates epithelial bladder cells speed and direction of movement during cell migration and tumor cell invasion. Inhibits Akt pro-survival-induced kinase activity. Mediates Rho protein-induced transcriptional activation via the c-fos serum response factor (SRF). Phosphorylates HCV NS5B leading to stimulation of HCV RNA replication.
Product Categories/Family for anti-PKN2 antibody	Signal Transduction

NCBI AND UNIPROT INFORMATION

NCBI GI #	1002635082
NCBI GeneID	5586
NCBI Accession #	NP_001307636.1
NCBI GenBank Nucleotide #	NM_001320707.1
UniProt Accession #	Q16513; Q08AF4; Q9H1W4; B4DQ21; B4DTP5; B4DVG1; D3DT24
Molecular Weight	75,171 Da
NCBI Official Full Name	serine/threonine-protein kinase N2 isoform 2
NCBI Official Synonym Full Names	protein kinase N2
NCBI Official Symbol	PKN2
NCBI Official Synonym Symbols	PAK2; PRK2; STK7; Pak-2; PRKCL2; PRO2042
NCBI Protein Information	serine/threonine-protein kinase N2
UniProt Protein Name	Serine/threonine-protein kinase N2
UniProt Gene Name	PKN2
UniProt Synonym Gene Names	PRK2; PRKCL2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://www.aaabiotech.com/polyclonal/pkn2-antibody-immunofluorescence-immunohistochemistry-elisa/233727>